

## Novelty-Facilitated Extinction: Providing a Novel Outcome in Place of an Expected Threat Diminishes Recovery of Defensive Responses

Joseph E. Dunsmoor, Vinn D. Campese, Ahmet O. Ceceli, Joseph E. LeDoux, and Elizabeth A. Phelps

### ABSTRACT

**BACKGROUND:** Experimental extinction serves as a model for psychiatric treatments based on associative learning. However, the effects of extinction are often transient, as evidenced by postextinction return of defensive behaviors. From a therapeutic perspective, an inherent problem with extinction may be that mere omission of threat is not sufficient to reduce future threat uncertainty. The current study tested an augmented form of extinction that replaced, rather than merely omitted, expected threat outcomes with novel nontreat outcomes, with the goal of reducing postextinction return of defensive behaviors.

**METHODS:** Thirty-two healthy male Sprague-Dawley rats and 47 human adults underwent threat conditioning to a conditioned stimulus paired with an electrical shock. Subjects then underwent a standard extinction protocol with shock omitted or an augmented extinction protocol wherein the shock was replaced by a surprising tone. Tests of postextinction recovery occurred 24 hours later in the absence of the tone.

**RESULTS:** Replacing the shock with a novel nontreat outcome, as compared with shock omission, reduced postextinction recovery (freezing in rats and anticipatory skin conductance responses in humans) when tested 24 hours later. Self-reported intolerance of uncertainty was positively correlated with recovery following standard extinction in humans, providing new evidence that postextinction recovery is related to sensitivity to future threat uncertainty.

**CONCLUSIONS:** These findings provide cross-species evidence of a novel strategy to enhance extinction that may have broad implications for how to override associative learning that has become maladaptive and offer a simple technique that could be straightforwardly adapted and implemented in clinical situations.

**Keywords:** Anxiety, Arousal, Electrodermal, Fear, Pavlovian, Regulation

<http://dx.doi.org/10.1016/j.biopsych.2014.12.008>

Without effort or intention, individuals retain information associated with highly aversive experiences. In extreme cases, this leads to persistent intrusive memories and unwanted physiological responses to cues associated with the event. Investigations on how to mitigate the psychological and physiological impact of negative events borrows heavily from research on extinction of classical threat (fear) conditioning, whereby the omission of an aversive event reduces defensive behaviors, like increases in sympathetic arousal and freezing. Laboratory research routinely shows, however, that defensive behaviors return postextinction (1–3). That learned defensive behaviors return even in laboratory experiments with healthy subjects reveals that threat conditioning is a powerful form of learning and that extinction is a rather unsatisfactory way to acquire safety. As the principles of extinction continue to serve as a model for clinical treatments like exposure therapy (4,5), there is strong motivation to discover innovative behavioral techniques to prevent postextinction return of defensive behaviors.

The weakness of extinction may be owed to a number of evolutionary and environmental factors (6,7). One limiting factor is that extinction does not eliminate the association between a conditioned stimulus (CS) and an unconditioned stimulus (US). Instead, extinction is a form of retroactive interference in which the new safety association competes for expression against the original threat association (6,8). This secondary association is fragile, as evidenced by the return of conditioned responses (CR) following the passage of time (spontaneous recovery), following presentation of the US (reinstatement), or when cues are encountered outside the extinction context (renewal) [see (9) for review]. Preventing the return of CRs remains a challenge (10). Models that conceptualize extinction as another form of associative learning propose that animals are provoked to learn by the surprising absence of the US (8,11). However, omission does not guarantee an effective learning signal. This is especially true in relapse following exposure treatment, where highly feared outcomes are repeatedly disconfirmed but continue to be expected following treatment, e.g., a persistent fear of dying in

an airplane crash after exposure therapy for fear of flying. In other words, the mere absence of the US is not sufficient to provoke a durable safety memory.

Another limit to extinction is that US omission can render the meaning of the CS ambiguous (12). That is, the CS takes on an additional meaning where it no longer predicts the US (or predicts no US). Resolving this ambiguity tends to favor expression of the original memory, perhaps since this association was learned first and/or is simply more salient than the extinction memory. For example, expression of threat conditioning readily generalizes to related cues and across multiple contexts, whereas extinction tends to be confined to the extinguished cue in the extinction context (12–14). The unexpected absence of the US could also promote a general sense of future threat uncertainty. If so, then individuals who have difficulty coping with uncertainty may be especially prone to return of defensive behaviors following extinction.

In the present study, we reasoned that extinction could be straightforwardly and effectively augmented if the aversive US was instead replaced by a surprising nonthreat outcome. Here, an expected electric shock was replaced by a novel and affectively neutral tone. We predicted that replacing shock with a novel outcome would reduce postextinction spontaneous recovery more effectively than shock omission alone for two principal reasons. First, unlike the mere absence of the shock, a novel outcome should be more effective at generating a mismatch between the predicted and received outcome, therefore signaling a clear change in the environment to promote the acquisition of new learning (8,15). Second, a novel perceptible outcome should help resolve some ambiguity generated by shock omission by providing a more substantive alternative association for the CS than no shock.

It is important to note up front certain methodological similarities to another form of outcome interference: counterconditioning. In counterconditioning, the outcome switches between opposing reinforcement qualities (appetitive and aversive) and thus opposing behavioral responses (approach and avoidance). Counterconditioning serves as a model for some behavior therapies (16,17), but contemporary research on aversive-to-appetitive counterconditioning in humans is scant (18,19). However, there is ample evidence in rats showing strong return of defensive CRs following counterconditioning (20–25). Unlike counterconditioning, the current protocol replaces the US with a surprising stimulus that does not reinforce any overt behavior.

The effect of this modified extinction protocol was assessed 24 hours after extinction in rats (experiment 1) and humans (experiment 2), with the prediction that a novel nonthreat outcome would diminish spontaneous recovery. We additionally explored for the first time whether individual differences in self-reported intolerance of uncertainty predicts postextinction recovery in humans.

## METHODS AND MATERIALS


### Experiment 1

**Animal Subjects.** Thirty-two male Sprague-Dawley rats (Hilltop Lab Animals, Scottsdale, Pennsylvania) were used in experiment 1. Further details regarding the animal subjects

and threat conditioning procedures are included in [Supplement 1](#).

**Rodent Threat Conditioning Procedures.** Two contexts were created (A and B) and made distinct with visual, tactile, and olfactory cues. Subjects were placed in context A on the first day of the study ([Figure 1A](#)), wherein the front panel light was illuminated for 30 seconds and co-terminated in the delivery of a 1-second .7 mA footshock on 100% of trials. The acquisition session included five trials spaced apart by an approximate 180-second intertrial interval. Two minutes following the last trial, subjects were removed from the chambers and returned to the colony for the remainder of the day.

Subjects were randomly assigned to one of two groups that differed only in regard to the type of extinction training: extinction through shock omission (EXT) or extinction modified by a surprising and novel nonthreat outcome in place of the shock, a procedure we refer to as novelty-facilitated extinction (NFE). Rats in both groups were placed in context B on day 2 of the study. Subjects in the EXT group received only the 30-second light presentation. For NFE subjects, the final 10 seconds of the light presentation were accompanied by a 5-kHz tone. This session included 30 trials separated by 15



**Figure 1.** (A) Procedure for experiment 1 in rats. Two groups underwent identical differential threat conditioning to initiate freezing to a conditioned stimulus (CS) paired with an electric shock unconditioned stimulus (US). Twenty-four hours later in a new context, one group underwent standard extinction in which the US was omitted on CS trials, while another group underwent a modified extinction procedure in which the US was replaced by a novel nonaversive tone. Spontaneous recovery was tested for both groups 24 hours later in the extinction context. (B) Freezing during extinction in the two groups was equivalent. (C) The novelty-facilitated extinction (NFE) group showed significantly less evidence of recovery 24 hours later than the standard extinction group. \*\* $p < .01$ ; error bars reflect standard error. CS duration was 30 seconds. EXT, extinction through shock omission.

seconds with an additional 2 minutes following the final trial. No shocks were delivered during this session. The day after extinction, all subjects were returned to context B for a test of extinction retention to the extinguished light CS alone (five trials). Exploratory tests of conditioned inhibition and contextual renewal were then conducted in the same subjects to investigate the influence of NFE on response inhibition. Data from these tests were mixed and are not included here to ease the interpretation of the spontaneous recovery data.


**Rodent Behavior Analysis.** A rater blind to experimental conditions manually scored freezing behavior. Freezing is presented as a percentage of the total CS duration. Inspection of acquisition indicated successful acquisition, i.e., all subjects demonstrated shock reactivity and high rates of freezing (data not presented). Data from extinction were analyzed with split-plot repeated measures analysis of variance procedures using extinction-block (average of five CS trials) and group (EXT, NFE) as within and between subject factors, respectively. Planned *t* tests were used to directly compare freezing between groups during extinction and recovery tests and were considered significant at  $\alpha < .05$ , two-tailed.

## Experiment 2

**Human Participants.** All participants gave written informed consent approved by the University Committee on Activities Involving Human Subjects at New York University. Sixty healthy volunteers participated in experiment 2. Thirteen participants were excluded from the final analysis due to technical problems ( $n = 2$ ), failure to attend to the task ( $n = 2$ ),

or failure of threat conditioning as defined by a positive difference in mean skin conductance responses (SCR) to the CS that predicted shock (CS+) versus a control stimulus that was never paired with shock (CS-) in the second half of threat conditioning ( $n = 9$ ). Removing these subjects (nonlearners) is justified as this investigation is focused on recovery from extinction, which is predicated on participants having evinced acquisition, and is standard practice in studies investigating extinction manipulations [e.g., (26–28)]. The final sample included 47 participants (25 female subjects; mean  $\pm$  SD:  $24.06 \pm 6.79$ ). A third group was run to investigate the effects of avoidance responses on postextinction recovery, but data from this group are not included here. Following informed consent, participants completed the State and Trait Anxiety Inventory (29) and the Intolerance of Uncertainty Scale (IUS) (30) and then randomly assigned to the EXT group ( $n = 23$ ; 13 female subjects) or the NFE group ( $n = 24$ , 12 female subjects).

**Human Threat Conditioning Procedures.** Experiment 2 occurred over 2 days separated by 24 hours (Figure 2A). Conditioned stimuli were two angry faces (31) presented for 6 seconds that signaled the presence (CS+) or absence (CS-) of an aversive electric shock to the right wrist during threat conditioning. Faces serving as CS+ and CS- were counterbalanced. Each trial was separated by an intertrial interval of 12 seconds. Subjects in all experimental groups wore headphones (Sennheiser HD-280 PRO; Sennheiser, Wedemark, Germany), which they were told were needed to block out background noise. This provided a convenient cover story for



**Figure 2.** (A) Procedure for experiment 2 in humans. Two groups underwent identical differential threat conditioning to initiate conditioned skin conductance responses (SCRs) to a conditioned stimulus (CS+) paired with an electric shock unconditioned stimulus (US). An unpaired CS (CS-) served as a within-subjects control condition. One group then underwent standard extinction in which the US was omitted on CS+ trials, while another group underwent a modified extinction procedure in which the US was replaced by a novel nonaversive tone. Spontaneous recovery of SCRs to the CS+ versus CS- was tested 24 hours later. (B) SCR results showed equivalent acquisition (CS+ minus CS-) and extinction on day 1. The novelty-facilitated extinction (NFE) group showed no evidence of recovery 24 hours later, whereas the standard extinction group showed robust recovery. \* $p < .05$ , \*\* $p < .01$ ; error bars reflect standard error. EXT, extinction through shock omission; IUS, Intolerance for Uncertainty Scale; n.s., not significant;  $\mu$ S, square root transformed SCRs, in microsiemens.

the NFE group who later received surprising tones through the headphones during the augmented extinction session.

Conditioning occurred over two runs that each included 8 presentations of the CS+ and CS- (32 total trials). An additional eight CS+ trials (four in each run) co-terminated with the US (33% reinforcement rate). CS+ trials paired with shock were not included in the data analysis to mitigate any potential confound in the psychophysiological analysis introduced by the shock; thus, the number of CS+ and CS- trials in the analysis were equal. A short habituation phase preceded acquisition to reduce initial orienting responses, which contained four trials each of the CS+ and CS- (data not reported).

Following acquisition, subjects were randomly assigned to group EXT or group NFE in which a low-volume (>60 decibel) 440-Hz tone was unexpectedly presented and co-terminated with each CS+ trial in place of the electric shock for 1.5 seconds. In both groups, extinction occurred over two runs that included 8 presentations each of the CS+ and CS- (32 total trials) in the absence of the US. Subjects returned 24 hours later, at which time the shock electrodes were reattached and the intensity level was set to the level reached the previous day. The shock intensity was not recalibrated so as to avoid reinstatement before the spontaneous recovery test. The recovery test included the CS+ and CS- (10 trials of each) in the absence of shock. Additional details on the electrical shock, collection of skin conductance responses, and an exploratory test of reinstatement are included in [Supplement 1](#).

**Human Psychophysiology Data Analysis.** SCRs were analyzed separately for early and late trials, as differences in responding between the CS+ and CS- tended to vary over the course of training [see also (32)]. The late phase of threat conditioning was defined as the mean of the last eight CS+ and eight CS- trials. The late phase of extinction was defined as the last three CS+ and three CS- trials. As the late phases of acquisition and extinction are the most relevant to show learning effects, only the late phases for acquisition and extinction are reported. Additionally, as prior research has shown that extinction manipulations tend to affect the earliest presentations of the CS+ versus the CS- in recovery tests (33), the early phase of spontaneous recovery was defined as the mean of the first three CS+ trials and first three CS- trials. The late phase of extinction retention was defined as the mean of the last three trials of the CS+ and three CS-. Results from each phase were analyzed by repeated measures analysis of variance with stimulus (CS+, CS-) as a within-subjects factor and group as a between-subjects factor, considered significant at  $\alpha < .05$ , two-tailed. We provide a complementary analysis of recovery that accounts for individual differences in acquisition, adapted from Milad *et al.* (34). We calculated a spontaneous recovery index by dividing each subject's mean SCRs to the CS+ during early recovery by the largest SCR to the CS+ during acquisition on day 1. This value provides a percent of recovery as a function of initial acquisition, with higher values indicating stronger recovery as a function of initial conditioning, whereas lower values indicate lower recovery as a function of initial conditioning.

## RESULTS

### Experiment 1: Novelty-Facilitated Extinction Versus Standard Extinction in Rats

**Extinction.** Freezing diminished over the course of extinction training in both groups ([Figure 1B](#)). There was a significant main effect for trial block (average of five trials;  $F_{5,15} = 9.16$ ,  $p < .001$ ) and no significant effect of group ( $p = .21$ ); thus, extinction proceeded comparably for both groups. Furthermore, there was no difference in freezing in the last five trials of extinction between the two groups ( $t_{30} = 1.28$ ,  $p = .21$ ), demonstrating that they were at an equivalent level of freezing by the end of extinction.

**24-Hour Extinction Retention Test.** Extinction retention tests showed that subjects in the EXT group froze significantly more than subjects in the NFE group ( $t_{30} = 2.70$ ,  $p = .01$ ,  $d = .87$ ), indicating that the NFE protocol diminished recovery of defensive responses compared with mere omission of shock (i.e., standard extinction) in rats ([Figure 1C](#)).

### Experiment 2: Novelty-Facilitated Extinction Versus Standard Extinction in Humans

**Acquisition.** SCRs from late threat conditioning showed equivalent levels of acquisition between groups ([Figure 2B](#)). There was a main effect of CS ( $F_{1,45} = 63.11$ ,  $p < .001$ ,  $\eta^2_p = .58$ ) but no group interaction ( $p = .99$ ). Planned *t* tests on differential SCRs (CS+ minus CS-) confirmed acquisition in group EXT ( $t_{22} = 5.26$ ,  $p < .001$ ) and group NFE ( $t_{23} = 6.02$ ,  $p < .001$ ), with no difference between groups ( $p = .93$ ).

**Extinction.** By late extinction, there was no effect of CS ( $p = .13$ ) and no group interaction ( $p = .15$ ). As confirmation of extinction, *t* tests performed on the late extinction SCR difference scores were not significant for either group ( $p > .05$ ).


**24-Hour Spontaneous Recovery Test.** The primary prediction in this experiment was that replacing the US with a novel nonthreat outcome would reduce postextinction recovery more effectively than merely omitting the US altogether. The early phase of spontaneous recovery was characterized by a main effect of CS ( $F_{1,45} = 18.72$ ,  $p < .001$ ,  $\eta^2_p = .29$ ) as well as a CS  $\times$  group interaction ( $F_{1,45} = 5.94$ ,  $p = .019$ ,  $\eta^2_p = .12$ ). Planned *t* tests revealed greater differential SCRs (CS+ minus CS-) in the EXT relative to the NFE group ( $t_{45} = 2.43$ ,  $p = .019$ ,  $d = .71$ ). Consistent with the CS  $\times$  group interaction, paired *t* tests showed no SCR difference between the CS+ (mean  $\pm$  standard error:  $.41 \pm .07$ ) and CS- ( $.33 \pm .08$ ) in the NFE group ( $t_{23} = 1.35$ ,  $p = .19$ ) but significantly greater SCRs to CS+ ( $.74 \pm .09$ ) than to the CS- ( $.43 \pm .06$ ) in the EXT group ( $t_{22} = 4.73$ ,  $p < .001$ ). These effects persisted into late recovery, where there remained a main effect of CS type ( $F_{1,45} = 7.33$ ,  $p = .01$ ,  $\eta^2_p = .140$ ) and a CS  $\times$  group interaction ( $F_{1,45} = 6.96$ ,  $p = .01$ ,  $\eta^2_p = .13$ ). Again, this effect was driven by a greater SCR difference score in the EXT relative to NFE group ( $t_{45} = 2.63$ ,  $p = .011$ ,  $d = .77$ ). Consistent with analysis

on early recovery SCRs, the spontaneous recovery index also showed greater recovery in the EXT ( $57.28\% \pm 5.96\%$ ) relative to the NFE ( $31.23\% \pm 5.34\%$ ) group ( $t_{45} = 3.37, p < .01$ ) (Figure 3A). Overall, these results provide evidence that a novel nonthreat outcome during extinction reduced postextinction recovery relative to standard extinction in which the US was merely omitted.

**Correlations between Spontaneous Recovery and Intolerance of Uncertainty.** One potential explanation for the fragility of standard extinction is that the omission of the US could in some individuals trigger a general sense of future threat uncertainty (35). That is, rather than instilling a sense of safety, omission renders the probability of shock less predictable in the long term. This uncertainty could interfere with the formation of a persistent extinction memory and lead to threat recovery when the CS is re-encountered. One approach to test this hypothesis is to examine whether threat recovery following extinction is related to the individual subject's inability to cope with ambiguity and uncertainty, as measured by subjective report on the IUS. Intolerance of uncertainty (30,36) is characteristic of individuals who tend to worry excessively (37), a trait characterized by interpreting ambiguous events as threatening and overestimating risk (38). There was a positive correlation between SCRs during early spontaneous recovery and IUS scores in the EXT group ( $R^2_{21} = .23, p = .022$ ) but not in the NFE group ( $p = .89$ ) (Figure 2C). The correlation in the EXT group remained significant after controlling for trait anxiety levels,  $R^2_{20} = .21, p = .034$ . A correlation with IUS scores was also observed using the spontaneous recovery index ( $R^2_{21} = .29, p = .008$ ) but not in the NFE group ( $p = .89$ ) (Figure 3B). This association also held after controlling for trait anxiety levels ( $R^2_{20} = .24, p = .019$ ), and a test of the difference between correlation coefficients revealed a significant difference between the EXT and NFE group ( $p = .04$ ).

## DISCUSSION

Results show that replacing an aversive shock with a novel nonaversive outcome was more effective at diminishing recovery of conditioned responses in rats and humans than



extinction that relied on omission alone. Effective extinction learning necessitates that the initial CS-US association is updated and modified by new information. Surprise encourages this new learning by signaling change (15). Thus, when extinction fails to yield a persistent memory, it could be due to the ineffectiveness of omission alone signaling a clear change in the meaning of the CS. Consequently, reductions in defensive responses seen over the course of standard extinction training do not necessarily reflect the formation of a durable safety memory (39,40).

In conditioning research, two heavily investigated forms of retroactive interference are extinction and counterconditioning (41). Both types of new learning interfere with the original CS-US association, but the original CR is prone to recovery (20). Thus, the results obtained here are in contrast to those predicted from a counterconditioning protocol. Theoretical accounts of counterconditioning have proposed that the effect relies on competing motivational states and competing expectations between different reinforcements with incompatible responses (17). In the case of aversive-to-appetitive counterconditioning, reinforcement by a new appetitive outcome upholds the CS as an important cue in the environment that commands a response—the manner determined by which memory representation is retrieved at the time, approach, or withdrawal. Indeed, counterconditioning is also referred to as cross-motivational transfer in the learning literature [e.g., (22)], implying the goal of the technique is to transfer responses from one set of motivated behaviors to an opposing set of motivated behaviors. Here, we incorporated an outcome interference design that did not reinforce a behavioral response. Consequently, subjects may have reinterpreted the CS as behaviorally irrelevant through association with a novel neutral outcome. If the goal of new learning is to reduce the relevance of threat conditioned cues altogether, then counterconditioning may not be the most appropriate technique.

The technique used here can also be distinguished from response inhibition techniques that focus on reducing the aversiveness of the US itself, including US habituation (42) and US revaluation (43). In the present study, the meaning or intensity of the US was not targeted. Rather, we focused on

**Figure 3.** (A) A complementary index of recovery was created for each subject by dividing mean skin conductance responses to the CS+ during early recovery by the subject's largest skin conductance response to the CS+ during acquisition on day 1. This value was multiplied by 100, providing an index of recovery as a function of initial responding for each subject. The standard extinction group showed significantly greater recovery by this index than the novelty-facilitated extinction (NFE) group. (B) Index of 24-hour spontaneous recovery correlated with self-reported intolerance of uncertainty (Intolerance of Uncertainty Scale [IUS]) in the standard extinction group. These correlations were selective to the standard extinction group. \*\* $p < .01$ ; error bars reflect standard error. CS+, conditioned stimulus signaling shock; EXT, extinction through shock omission.

the uncertainty inherent to omission of the US during extinction by replacing the US with a novel outcome.

The cross-species design of this study provides generalizability across different levels of behavior, from freezing in rats to autonomic arousal in humans. It has been suggested that SCRs are a cognitive expression of conditioned learning (44). That the NFE procedure was effective in rats, however, provides evidence that these effects are not due to some cognitive factor specific to humans. Importantly, in humans we were able to assess subjective ratings of intolerance of uncertainty to test the hypothesis that recovery following standard extinction is related to individual differences in intolerance of uncertainty, affording a different level of analysis than is possible from an animal model. Positive findings of the NFE technique in rats also open the possibility for more extensive future research on the neurophysiology underlying this effect than is possible using humans, with a likely focus on regions implicated in novelty-enhanced learning, including the hippocampus and striatum (45).

The inclusion of human subjects also provides a closer step toward understanding the clinical implications of the present results. The NFE technique may be appropriate for therapeutic purposes when trying to diminish the perceived significance of situations or stimuli inappropriately associated with danger. In other words, an association with an unremarkable outcome could restore the neutrality of everyday events and objects. Clinical applications could include augmenting exposure therapy to increase surprise using novel but unremarkable outcomes, in line with clinical models that highlight the critical role for belief disconfirmation for the success of exposure treatments (46,47). For example, in exposure treatments, it may be more beneficial to have negative beliefs disconfirmed by experiencing an unexpectedly mundane event, rather than simply experiencing the absence of a negative event. These findings lend support to the idea that updating beliefs for negative outcomes may occur more efficiently through this sort of surprise and also fit with associative learning models that emphasize the importance of surprise on updating previously held knowledge (8,15).

While much research describes postextinction return of defensive responses as an unwanted consequence of unsuccessful extinction, it is important to point out that the failure to completely override threat associations serves an adaptive purpose. That is, it is advantageous to maintain a memory that certain cues signal danger despite some evidence to the contrary, because even one encounter with a threat (e.g., a predator) threatens survival. In other words, fully effective extinction could lead to the incorrect conclusion that a cue is entirely safe, with disastrous results. Extinction learning provides animals with a repertoire of flexible behaviors that are (ideally) generated under the appropriate circumstances. To utilize the principles of extinction as a therapeutic tool, one must overcome the limits to extinction, which are inherently beneficial under survival circumstances but maladaptive in pathological anxiety. Learning through surprise that a CS with a prior history of predicting danger now predicts something trivial may help overcome these limits.

We note some limitations to the present studies. First, future research using the NFE paradigm in clinical populations is

needed to generalize these results to anxiety patients. Future experiments will also be needed to determine how NFE targets other postextinction phenomena, including contextual renewal and reinstatement. Future studies in humans should consider incorporating immersive virtual reality to emulate characteristics of real world environments (48). Finally, an intriguing possibility is that NFE transforms the CS+ into a conditioned inhibitor (i.e., safety signal). Such an effect would be a powerful demonstration that the NFE technique is far more effective than standard extinction, which does not result in the CS taking on the properties of a safety signal [see also (49)].

Finally, these results provide possible explanations as to why extinction often fails to instill persistent memories, as individuals who express an inability to cope with ambiguity and uncertainty were more susceptible to threat recovery following standard extinction. The novelty-facilitated extinction technique presented here provides a straightforward technique to help mitigate uncertainty inherent to extinction procedures and may be especially beneficial to reduce relapse in individuals with high intolerance to uncertainty. Insofar as the principles of threat conditioning and extinction have been successfully extended to understand and treat clinical anxiety (5), this finding offers new insights into the types of individuals who may be more likely to relapse following exposure therapy and establishes a strategy that is simple to implement and may provide more effective clinical outcomes than strategies based on merely omitting aversive events.

## ACKNOWLEDGMENTS AND DISCLOSURES

This study was supported by National Institutes of Health RO1 MH097085, National Institutes of Health Grant M0H38774, and National Institute of Mental Health Training Award in Systems and Integrative Neuroscience T32 MH019524.

All authors report no biomedical financial interests or potential conflicts of interest.

## ARTICLE INFORMATION

From the Psychology Department and Center for Neural Sciences (JED, VDC, AOC, JEL, EAP), New York University, New York; and Emotional Brain Institute (JEL, EAP), Nathan Kline Institute for Psychiatric Research, Orangeburg, New York.

Address correspondence to Elizabeth A. Phelps, Ph.D., New York University, Psychology Department and Center for Neural Sciences, Center for Neural Science, 6 Washington Place Room 890, New York, NY 10003; E-mail: liz.phelps@nyu.edu.

Received Aug 21, 2014; revised Nov 7, 2014; accepted Dec 3, 2014.

Supplementary material cited in this article is available online at <http://dx.doi.org/10.1016/j.biopsych.2014.12.008>.

## REFERENCES

1. Vervliet B, Craske MG, Hermans D (2013): Fear extinction and relapse: State of the art. *Annu Rev Clin Psychol* 9:215–248.
2. Bouton ME (2002): Context, ambiguity, and unlearning: Sources of relapse after behavioral extinction. *Biol Psychiatry* 52:976–986.
3. Rescorla RA (2004): Spontaneous recovery. *Learn Mem* 11:501–509.
4. Foa EB, Steketee G, Rothbaum BO (1989): Behavioral cognitive conceptualizations of post-traumatic stress disorder. *Behav Ther* 20: 155–176.
5. Milad MR, Quirk GJ (2012): Fear extinction as a model for translational neuroscience: Ten years of progress. *Annu Rev Psychol* 63:129–151.

6. Bouton ME (1993): Context, time, and memory retrieval in the interference paradigms of Pavlovian learning. *Psychol Bull* 114:80–99.
7. Larrauri JA, Schmajuk NA (2008): Attentional, associative, and configural mechanisms in extinction. *Psychol Rev* 115:640–676.
8. Pearce JM, Hall G (1980): A model for Pavlovian learning: Variations in the effectiveness of conditioned but not of unconditioned stimuli. *Psychol Rev* 87:532–552.
9. McConnell BL, Miller RR (2014): Associative accounts of recovery-from-extinction effects. *Learn Motiv* 46:1–15.
10. Fitzgerald PJ, Seemann JR, Maren S (2014): Can fear extinction be enhanced? A review of pharmacological and behavioral findings. *Brain Res Bull* 105C:46–60.
11. Rescorla RA, Wagner AR (1972): A Theory of Pavlovian Conditioning: Variations in the Effectiveness of Reinforcement and Nonreinforcement. New York: Appleton-Century-Crofts.
12. Bouton ME (2004): Context and behavioral processes in extinction. *Learn Mem* 11:485–494.
13. Vervliet B, Vansteenwegen D, Baeyens F, Hermans D, Eelen P (2005): Return of fear in a human differential conditioning paradigm caused by a stimulus change after extinction. *Behav Res Ther* 43:357–371.
14. Vervoort E, Vervliet B, Bennett M, Baeyens F (2014): Generalization of human fear acquisition and extinction within a novel arbitrary stimulus category. *PLoS One* 9:e96569.
15. Courville AC, Daw ND, Touretzky DS (2006): Bayesian theories of conditioning in a changing world. *Trends Cogn Sci* 10:294–300.
16. Paunovic N (2003): Prolonged exposure counterconditioning as a treatment for chronic posttraumatic stress disorder. *J Anxiety Disord* 17:479–499.
17. Wolpe J (1952): Experimental neuroses as learned behaviour. *Br J Psychol Gen Sect* 43:243–268.
18. Bulgaria L, Bach DR, Wittmann BC (2014): Prior fear conditioning and reward learning interact in fear and reward networks. *Front Behav Neurosci* 8:67.
19. Raes AK, De Raedt R (2012): The effect of counterconditioning on evaluative responses and harm expectancy in a fear conditioning paradigm. *Behav Ther* 43:757–767.
20. Delamater AR (1996): Effects of several extinction treatments upon the integrity of Pavlovian stimulus-outcome associations. *Anim Learn Behav* 24:437–449.
21. Brooks DC, Hale B, Nelson JB, Bouton ME (1995): Reinstatement after counterconditioning. *Anim Learn Behav* 23:383–390.
22. Bouton ME, Peck CA (1992): Spontaneous recovery in cross-motivational transfer (counterconditioning). *Anim Learn Behav* 20: 313–321.
23. Peck CA, Bouton ME (1990): Context and performance in aversive-to-appetitive and appetitive-to-aversive transfer. *Learn Motiv* 21:1–31.
24. Thomas BL, Cutler M, Novak C (2012): A modified counterconditioning procedure prevents the renewal of conditioned fear in rats. *Learn Motiv* 43:24–34.
25. Tunstall BJ, Verendeev A, Kearns DN (2012): A comparison of therapies for the treatment of drug cues: Counterconditioning vs. extinction in male rats. *Exp Clin Psychopharmacol* 20:447–453.
26. Kindt M, Soeter M (2013): Reconsolidation in a human fear conditioning study: A test of extinction as updating mechanism. *Biol Psychol* 92:43–50.
27. Kalisch R, Korenfeld E, Stephan KE, Weiskopf N, Seymour B, Dolan RJ (2006): Context-dependent human extinction memory is mediated by a ventromedial prefrontal and hippocampal network. *J Neurosci* 26: 9503–9511.
28. Milad MR, Orr SP, Pitman RK, Rauch SL (2005): Context modulation of memory for fear extinction in humans. *Psychophysiology* 42: 456–464.
29. Spielberger CD (1983): Manual for the State-Trait Anxiety Inventory. Palo Alto, CA: Consulting Psychologists Press.
30. Buhr K, Dugas MJ (2002): The intolerance of uncertainty scale: Psychometric properties of the English version. *Behav Res Ther* 40: 931–945.
31. Ekman P, Friesen WV (1976): Measuring facial movement. *Environmental Psychology and Nonverbal Behavior* 1:56–75.
32. Rabinak CA, Angstadt M, Sripada CS, Abelson JL, Liberzon I, Milad MR, Phan KL (2013): Cannabinoid facilitation of fear extinction memory recall in humans. *Neuropharmacology* 64:396–402.
33. Schiller D, Monfils MH, Raio CM, Johnson DC, LeDoux JE, Phelps EA (2010): Preventing the return of fear in humans using reconsolidation update mechanisms. *Nature* 463:49–53.
34. Milad MR, Wright CI, Orr SP, Pitman RK, Quirk GJ, Rauch SL (2007): Recall of fear extinction in humans activates the ventromedial prefrontal cortex and hippocampus in concert. *Biol Psychiatry* 62: 446–454.
35. Grue DW, Nitschke JB (2013): Uncertainty and anticipation in anxiety: An integrated neurobiological and psychological perspective. *Nat Rev Neurosci* 14:488–501.
36. Freeston MH, Rhéaume J, Letarte H, Dugas MJ, Ladouceur R (1994): Why do people worry? *Pers Individ Dif* 17:791–802.
37. Dugas MJ, Gagnon F, Ladouceur R, Freeston MH (1998): Generalized anxiety disorder: A preliminary test of a conceptual model. *Behav Res Ther* 36:215–226.
38. Butler G, Mathews A (1987): Anticipatory anxiety and risk perception. *Cognit Ther Res* 11:551–565.
39. Craske MG, Kircanski K, Zelikowsky M, Mystkowski J, Chowdhury N, Baker A (2008): Optimizing inhibitory learning during exposure therapy. *Behav Res Ther* 46:5–27.
40. Rowe MK, Craske MG (1998): Effects of varied-stimulus exposure training on fear reduction and return of fear. *Behav Res Ther* 36:719–734.
41. Pavlov IP (1927): Conditioned Reflexes. London: Oxford University Press.
42. Storsve AB, McNally GP, Richardson R (2012): Renewal and reinstatement of the conditioned but not the unconditioned response following habituation of the unconditioned stimulus. *Behav Processes* 90: 58–65.
43. Dibbets P, Poort H, Arntz A (2012): Adding imagery rescripting during extinction leads to less ABA renewal. *J Behav Ther Exp Psychiatry* 43: 614–624.
44. Sevenster D, Beckers T, Kindt M (2012): Instructed extinction differentially affects the emotional and cognitive expression of associative fear memory. *Psychophysiology* 49:1426–1435.
45. Izquierdo LA, Barros DM, Medina JH, Izquierdo I (2000): Novelty enhances retrieval of one-trial avoidance learning in rats 1 or 31 days after training unless the hippocampus is inactivated by different receptor antagonists and enzyme inhibitors. *Behav Brain Res* 117: 215–220.
46. Salkovskis PM, Hackmann A, Wells A, Gelder MG, Clark DM (2007): Belief disconfirmation versus habituation approaches to situational exposure in panic disorder with agoraphobia: A pilot study. *Behav Res Ther* 45:877–885.
47. Craske MG, Treanor M, Conway CC, Zbozinek T, Vervliet B (2014): Maximizing exposure therapy: An inhibitory learning approach. *Behav Res Ther* 58C:10–23.
48. Dunsmoor JE, Ahs F, Zielinski DJ, LaBar KS (2014): Extinction in multiple virtual reality contexts diminishes fear reinstatement in humans. *Neurobiol Learn Mem* 113:157–164.
49. Leung HT, Reeks LM, Westbrook RF (2012): Two ways to deepen extinction and the difference between them. *J Exp Psychol Anim Behav Process* 38:394–406.