

NUMBERED KNOCKOUT PAPERS

1. Crabbe JC, Phillips TJ, Feller DJ, Hen R, Wenger CD, Lessov CN, Schafer GL, Elevated alcohol consumption in null mutant mice lacking 5-HT1B serotonin receptors, *Nat Genet*, 1996, 14:98-101.
2. Risinger FO, Bormann NM, Oakes RA, Reduced sensitivity to ethanol reward, but not ethanol aversion, in mice lacking 5-HT1B receptors, *Alcohol Clin Exp Res*, 1996, 20:1401-1405.
3. Miyakawa T, Yagi T, Kitazawa H, Yasuda M, Kawai N, Tsuboi K, Niki H, Fyn-kinase as a determinant of ethanol sensitivity: relation to NMDA-receptor function, *Science*, 1997, 278:698-701.
4. Homanics GE, Ferguson C, Quinlan JJ, Daggett J, Snyder K, Lagenaur C, Mi ZP, Wang XH, Grayson DR, Firestone LL, Gene knockout of the alpha6 subunit of the gamma-aminobutyric acid type A receptor: lack of effect on responses to ethanol, pentobarbital, and general anesthetics, *Mol Pharmacol*, 1997, 51:588-596.
5. Rubinstein M, Phillips TJ, Bunzow JR, Falzone TL, Dziewczapolski G, Zhang G, Fang Y, Larson JL, McDougall JA, Chester JA, Saez C, Pugsley TA, Gershanik O, Low MJ, Grandy DK, Mice lacking dopamine D4 receptors are supersensitive to ethanol, cocaine, and methamphetamine, *Cell*, 1997, 90:991-1001.
6. Timpl P, Spanagel R, Sillaber I, Kresse A, Reul JM, Stalla GK, Blanquet V, Steckler T, Holsboer F, Wurst W, Impaired stress response and reduced anxiety in mice lacking a functional corticotropin-releasing hormone receptor 1, *Nat Genet*, 1998, 19:162-166.
7. Quinlan JJ, Homanics GE, Firestone LL, Anesthesia sensitivity in mice that lack the beta3 subunit of the gamma-aminobutyric acid type A receptor, *Anesthesiology*, 1998, 88:775-780.
8. El Ghundi M, George SR, Drago J, Fletcher PJ, Fan T, Nguyen T, Liu C, Sibley DR, Westphal H, O'Dowd BF, Disruption of dopamine D1 receptor gene expression attenuates alcohol-seeking behavior, *Eur J Pharmacol*, 1998, 353:149-158.
9. Homanics GE, Le NQ, Kist F, Mihalek R, Hart AR, Quinlan JJ, Ethanol tolerance and withdrawal responses in GABA(A) receptor alpha 6 subunit null allele mice and in inbred C57BL/6J and strain 129/SvJ mice, *Alcohol Clin Exp Res*, 1998, 22:259-265.
10. Korpi ER, Koikkalainen P, Vekovischeva OY, Makela R, Kleinz R, Uusi-Oukari M, Wisden W, Cerebellar granule-cell-specific GABAA receptors attenuate benzodiazepine-induced ataxia: evidence from alpha 6-subunit-deficient mice, *Eur J Neurosci*, 1999, 11:233-240.

11. Homanics GE, Harrison NL, Quinlan JJ, Krasowski MD, Rick CE, de Blas AL, Mehta AK, Kist F, Mihalek RM, Aul JJ, Firestone LL, Normal electrophysiological and behavioral responses to ethanol in mice lacking the long splice variant of the gamma2 subunit of the gamma-aminobutyrate type A receptor, *Neuropharmacology*, 1999, 38:253-265.
12. Grisel JE, Mogil JS, Grahame NJ, Rubinstein M, Belknap JK, Crabbe JC, Low MJ, Ethanol oral self-administration is increased in mutant mice with decreased beta-endorphin expression, *Brain Res*, 1999, 835:62-67.
13. Risinger FO, Doan AM, Vickrey AC, Oral operant ethanol self-administration in 5-HT1b knockout mice, *Behav Brain Res*, 1999, 102:211-215.
14. Olive MF, Mehmert KK, Messing RO, Hodge CW, Reduced operant ethanol self-administration and in vivo mesolimbic dopamine responses to ethanol in PKCepsilon-deficient mice, *Eur J Neurosci*, 2000, 12:4131-4140.
15. Risinger FO, Freeman PA, Rubinstein M, Low MJ, Grandy DK, Lack of operant ethanol self-administration in dopamine D2 receptor knockout mice, *Psychopharmacology*, 2000, 152:343-350.
16. Cunningham CL, Howard MA, Gill SJ, Rubinstein M, Low MJ, Grandy DK, Ethanol-conditioned place preference is reduced in dopamine D2 receptor-deficient mice, *Pharmacol Biochem Behav*, 2000, 67:693-699.
17. Roberts AJ, McDonald JS, Heyser CJ, Kieffer BL, Matthes HW, Koob GF, Gold LH, mu-Opioid receptor knockout mice do not self-administer alcohol, *J Pharmacol Exp Ther*, 2000, 293:1002-1008.
18. Popova NK, Vishnivetskaya GB, Ivanova EA, Skrinskaya JA, Seif I, Altered behavior and alcohol tolerance in transgenic mice lacking MAO-A: a comparison with effects of MAO-A inhibitor clorgyline, *Pharmacol Biochem Behav*, 2000, 67:719-727.
19. Thiele TE, Miura GI, Marsh DJ, Bernstein IL, Palmiter RD, Neurobiological responses to ethanol in mutant mice lacking neuropeptide Y or the Y5 receptor, *Pharmacol Biochem Behav*, 2000, 67:683-691.
20. Weinshenker D, Rust NC, Miller NS, Palmiter RD, Ethanol-associated behaviors of mice lacking norepinephrine, *J Neurosci*, 2000, 20:3157-3164.
21. Thiele TE, Willis B, Stadler J, Reynolds JG, Bernstein IL, McKnight GS, High ethanol consumption and low sensitivity to ethanol-induced sedation in protein kinase A-mutant mice, *J Neurosci*, 2000, 20:RC75.

22. Bouwknecht JA, Hijzen TH, van der Gugten J, Maes RA, Hen R, Olivier B, Ethanol intake is not elevated in male 5-HT(1B) receptor knockout mice, *Eur J Pharmacol*, 2000, 403:95-98.
23. Bowers BJ, Elliott KJ, Wehner JM, Differential sensitivity to the anxiolytic effects of ethanol and flunitrazepam in PKCgamma null mutant mice, *Pharmacol Biochem Behav*, 2001, 69:99-110.
24. El Yacoubi M, Ledent C, Parmentier M, Daoust M, Costentin J, Vaugeois J, Absence of the adenosine A(2A) receptor or its chronic blockade decrease ethanol withdrawal-induced seizures in mice, *Neuropharmacology*, 2001, 40:424-432.
25. LaBuda CJ, Fuchs PN, The anxiolytic effect of acute ethanol or diazepam exposure is unaltered in mu-opioid receptor knockout mice, *Brain Res Bull*, 2001, 55:755-760.
26. Mihalek RM, Bowers BJ, Wehner JM, Kralic JE, VanDoren MJ, Morrow AL, Homanics GE, GABA(A)-receptor delta subunit knockout mice have multiple defects in behavioral responses to ethanol, *Alcohol Clin Exp Res*, 2001, 25:1708-1718.
27. Roberts AJ, Gold LH, Polis I, McDonald JS, Filliol D, Kieffer BL, Koob GF, Increased ethanol self-administration in delta-opioid receptor knockout mice, *Alcohol Clin Exp Res*, 2001, 25:1249-1256.
28. Otto C, Martin M, Wolfer DP, Lipp HP, Maldonado R, Schutz G, Altered emotional behavior in PACAP-type-I-receptor-deficient mice, *Mol Brain Res*, 2001, 92:78-84.
29. Risinger FO, Freeman PA, Greengard P, Fienberg AA, Motivational effects of ethanol in DARPP-32 knock-out mice, *J Neurosci*, 2001, 21:340-348.
30. Blednov YA, Stoffel M, Chang SR, Harris RA, Potassium channels as targets for ethanol: studies of G-protein-coupled inwardly rectifying potassium channel 2 (GIRK2) null mutant mice, *J Pharmacol Exp Ther*, 2001, 298:521-530.
31. Espinosa F, McMahon A, Chan E, Wang S, Ho CS, Heintz N, Joho RH, Alcohol hypersensitivity, increased locomotion, and spontaneous myoclonus in mice lacking the potassium channels Kv3.1 and Kv3.3, *J Neurosci*, 2001, 21:6657-6665.
32. Hall FS, Sora I, Uhl GR, Ethanol consumption and reward are decreased in mu-opiate receptor knockout mice, *Psychopharmacology*, 2001, 154:43-49.
33. Koenig HN, Olive MF, Ethanol consumption patterns and conditioned place preference in mice lacking preproenkephalin, *Neurosci Lett*, 2002, 325:75-78.

34. Horvath B, Spies C, Horvath G, Kox WJ, Miyamoto S, Barry S, Warden CH, Bechmann I, Diano S, Heemskerk J, Horvath TL, Uncoupling protein 2 (UCP2) lowers alcohol sensitivity and pain threshold, *Biochem Pharmacol*, 2002, 64:369-374.
35. Falzone TL, Gelman DM, Young JI, Grandy DK, Low MJ, Rubinstein M, Absence of dopamine D4 receptors results in enhanced reactivity to unconditioned, but not conditioned, fear, *Eur J Neurosci*, 2002, 15:158-164.
36. Becker A, Grecksch G, Kraus J, Loh HH, Schroeder H, Hollt V, Rewarding effects of ethanol and cocaine in mu opioid receptor-deficient mice, *Naunyn Schmiedebergs Arch Pharmacol*, 2002, 365:296-302.
37. Savelieva KV, Caudle WM, Findlay GS, Caron MG, Miller GW, Decreased ethanol preference and consumption in dopamine transporter female knock-out mice, *Alcohol Clin Exp Res*, 2002, 26:758-764.
38. Sillaber I, Rammes G, Zimmermann S, Mahal B, Zieglgansberger W, Wurst W, Holsboer F, Spanagel R, Enhanced and delayed stress-induced alcohol drinking in mice lacking functional CRH1 receptors, *Science*, 2002, 296:931-933.
39. Spanagel R, Siegmund S, Cowen M, Schroff KC, Schumann G, Fiserova M, Sillaber I, Wellek S, Singer M, Putzke J, The neuronal nitric oxide synthase gene is critically involved in neurobehavioral effects of alcohol, *J Neurosci*, 2002, 22:8676-8683.
40. Thiele TE, Koh MT, Pedrazzini T, Voluntary alcohol consumption is controlled via the neuropeptide Y Y1 receptor, *J Neurosci*, 2002, 22:RC208.
41. Choi DS, Wang D, Dadgar J, Chang WS, Messing RO, Conditional rescue of protein kinase C epsilon regulates ethanol preference and hypnotic sensitivity in adult mice, *J Neurosci*, 2002, 22:9905-9911.
42. Naassila M, Ledent C, Daoust M, Low ethanol sensitivity and increased ethanol consumption in mice lacking adenosine A2A receptors, *J Neurosci*, 2002, 22:10487-10493.
43. Narita M, Soma M, Tamaki H, Narita M, Suzuki T, Intensification of the development of ethanol dependence in mice lacking dopamine D(3) receptor, *Neurosci Lett*, 2002, 324:129-132.
44. Poncelet M, Maruani J, Calassi R, Soubrie P, Overeating, alcohol and sucrose consumption decrease in CB1 receptor deleted mice, *Neurosci Lett*, 2003, 343:216-218.
45. Palmer AA, Low MJ, Grandy DK, Phillips TJ, Effects of a *Drd2* deletion mutation on ethanol-induced locomotor stimulation and sensitization suggest a role for epistasis, *Behav Genet*, 2003, 33:311-324.

46. El Yacoubi M, Ledent C, Parmentier M, Costentin J, Vaugeois JM, Caffeine reduces hypnotic effects of alcohol through adenosine A2A receptor blockade, *Neuropharmacology*, 2003, 45:977-985.
47. Boyce-Rustay JM, Risinger FO, Dopamine D3 receptor knockout mice and the motivational effects of ethanol, *Pharmacol Biochem Behav*, 2003, 75:373-379.
- 48b. Blednov YA, Jung S, Alva H, Wallace D, Rosahl T, Whiting PJ, Harris RA Deletion of the alpha1 or beta2 subunit of GABAA receptors reduces actions of alcohol and other drugs, *J Pharmacol Exp Ther*, 2003, 304:30-36.
49. Hungund BL, Szakall I, Adam A, Basavarajappa BS, Vadasz C, Cannabinoid CB1 receptor knockout mice exhibit markedly reduced voluntary alcohol consumption and lack alcohol-induced dopamine release in the nucleus accumbens. *J Neurochem*, 2003, 84:698-704.
50. Kelai S, Aissi F, Lesch KP, Cohen-Salmon C, Hamon M, Lanfumey L, Alcohol intake after serotonin transporter inactivation in mice, *Alcohol Alcohol*, 2003, 38:386-389.
51. Hall FS, Sora I, Uhl GR, Sex-dependent modulation of ethanol consumption in vesicular monoamine transporter 2 (VMAT2) and dopamine transporter (DAT) knockout mice, *Neuropsychopharmacology*, 2003, 28:620-628.
52. Hill KG, Alva H, Blednov YA, Cunningham CL, Reduced ethanol-induced conditioned taste aversion and conditioned place preference in GIRK2 null mutant mice, *Psychopharmacology*, 2003, 169:108-114.
53. Kralic JE, Wheeler M, Renzi K, Ferguson C, O'Buckley TK, Grobin AC, Morrow AL, Homanics GE, Deletion of GABAA receptor alpha 1 subunit-containing receptors alters responses to ethanol and other anesthetics, *J Pharmacol Exp Ther*, 2003, 305:600-607.
54. Blednov YA, Stoffel M, Alva H, Harris RA, A pervasive mechanism for analgesia: activation of GIRK2 channels, *Proc Natl Acad Sci U S A*, 2003, 100:277-282.
- 55a. Blednov YA, Walker D, Alva H, Creech K, Findlay G, Harris RA, GABAA receptor alpha 1 and beta 2 subunit null mutant mice: behavioral responses to ethanol, *J Pharmacol Exp Ther*, 2003, 305:854-863.
56. Boehm SL 2nd, Peden L, Chang R, Harris RA, Blednov YA, Deletion of the fyn-kinase gene alters behavioral sensitivity to ethanol, *Alcohol Clin Exp Res*, 2003, 27:1033-1040.
57. Racz I, Bilkei-Gorzo A, Toth ZE, Michel K, Palkovits M, Zimmer A, A critical role for the cannabinoid CB1 receptors in alcohol dependence and stress-stimulated ethanol drinking, *J Neurosci*, 2003, 23:2453-2458.

58. Naassila M, Pierrefiche O, Ledent C, Daoust M, Decreased alcohol self-administration and increased alcohol sensitivity and withdrawal in CB1 receptor knockout mice, *Neuropharmacology*, 2004, 46:243-253.
59. Baskfield CY, Martin BR, Wiley JL, Differential effects of delta-9-tetrahydrocannabinol and methanandamide on operant behavior in CB1 knockout and wild type mice, *J Pharmacol Exp Ther*, 2004, 309:86-91.
60. Maul B, Siems WE, Hoehe MR, Grecksch G, Bader M, Walther T, Alcohol consumption is controlled by angiotensin II, *FASEB Journal*, 2001, 15:1640-1642.
61. Siems W, Maul B, Krause W, Gerard C, Hauser KF, Hersh LB, Fischer HS, Zernig G, Saria A, Neutral endopeptidase and alcohol consumption, experiments in neutral endopeptidase-deficient mice, *European Journal of Pharmacology*, 2000, 397:327-334.
62. Szumlinski KK, Toda S, Middaugh LD, Worley PF, Kalivas PW, Evidence for a relationship between Group 1 mGluR hypofunction and increased cocaine and ethanol sensitivity in Homer2 null mutant mice. In: B. Moghaddam and M.E. Wolf (Eds.), *Glutamate and Disorders of Cognition and Motivation* (series title: *Annals of the New York Academy of Sciences*, vol. 1003), New York Academy of Sciences, New York, 2003, pp. 468-471.
63. Ruan Y, Tecott L, Jiang MM, Jan LY, Jan YN, Ethanol hypersensitivity and olfactory discrimination defect in mice lacking a homolog of *Drosophila* neuralized, *Proceedings of the National Academy of Sciences USA*, 2001, 98:9907-9912.
64. Grahame NJ, Mosemiller AK, Low MJ, Froehlich JC, Naltrexone and alcohol drinking in mice lacking beta-endorphin by site-directed mutagenesis, *Pharmacology Biochemistry and Behavior*, 2000, 67:759-766.
65. Bechtholt AJ, Smith R, Raber J, Cunningham CL, Enhanced ethanol-, but not cocaine induced, conditioned place preference in *Apoe(-/-)* mice, *Pharmacology Biochemistry and Behavior*, 2004, 77:783-792.
66. Thiele TE, Naveilhan P, Ernfors P, Assessment of ethanol consumption and water drinking by NPY Y(2) receptor knockout mice, *Peptides*, 2004, 25:975-983.
67. Korkosz A, Kolomanska P, Kowalska K, Rogowski A, Radwanska K, Kaczmarek L, Mierzejewski P, Scinska A, Kostowski W, Bienkowski P, Dissociation of ethanol and saccharin preference in *fosB* knockout mice, *Physiol Behav*, 2004, 82:391-395.
68. Choi DS, Cascini MG, Mailliard W, Young H, Paredes P, McMahon T, Diamond I, Bonci A, Messing RO, The type 1 equilibrative nucleoside transporter regulates ethanol intoxication and preference, *Nat Neurosci*, 2004, 7:855-861.

69. Shelton KL, Dukat M, Allan AM, Effect of 5-HT₃ receptor over-expression on the discriminative stimulus effects of ethanol, *Alcohol Clin Exp Res*, 2004, 28:1161-1171.
70. Palmer AA, Sharpe AL, Burkhart-Kasch S, McKinnon CS, Coste SC, Stenzel-Poore MP, Phillips TJ, Corticotropin-releasing factor overexpression decreases ethanol drinking and increases sensitivity to the sedative effects of ethanol, *Psychopharmacology*, 2004, 176:386-397.
71. Pucilowski O, Ayensu WK, D'Ercole AJ, Insulin-like growth factor I expression alters acute sensitivity and tolerance to ethanol in transgenic mice, *Eur J Pharmacol*, 1996, 305:57-62.
72. Wick MJ, Radcliffe RA, Bowers BJ, Mascia MP, Luscher B, Harris RA, Wehner JM, Behavioural changes produced by transgenic overexpression of gamma₂L and gamma₂S subunits of the GABA_A receptor, *Eur J Neurosci*, 2000, 12:2634-2638.
73. Hu JH, Ma YH, Yang N, Mei ZT, Zhang MH, Fei J, Guo LH, Up-regulation of gamma-aminobutyric acid transporter I mediates ethanol sensitivity in mice, *Neuroscience*, 2004, 123:807-812.
74. Thiele TE, Marsh DJ, Ste Marie L, Bernstein IL, Palmiter RD, Ethanol consumption and resistance are inversely related to neuropeptide Y levels, *Nature*, 1998, 396:366-369.
75. Hilakivi-Clarke L, Goldberg R, Gonadal hormones and aggression-maintaining effect of alcohol in male transgenic transforming growth factor-alpha mice, *Alcohol Clin Exp Res*, 1995, 19:708-713.
76. Boehm SL 2nd, Peden L, Jennings AW, Kojima N, Harris RA, Blednov YA, Over-expression of the fyn-kinase gene reduces hypnotic sensitivity to ethanol in mice, *Neuroscience Letters*, 2004, 372:6-11.
77. de Fiebre NC, de Fiebre CM, alpha₇ Nicotinic acetylcholine receptor knockout selectively enhances ethanol-, but not beta-amyloid-induced neurotoxicity, *Neuroscience Letters*, 2005, 373:42-47.
78. Spanagel R, Pendyala G, Abarca C, Zghoul T, Sanchis-Segura C, Magnone MC, Lascorz J, Depner M, Holzberg D, Soyka M, Schreiber S, Matsuda F, Lathrop M, Schumann G, Albrecht U, The clock gene *Per2* influences the glutamatergic system and modulates alcohol consumption, *Nature Medicine*, 2005, 11:35-42.
79. Houchi H, Babovic D, Pierrefiche O, Ledent C, Daoust M, Naassila M, CB(1) receptor knockout mice display reduced ethanol-induced conditioned place preference and increased striatal dopamine D₂ receptors, *Neuropsychopharmacology*, 2005, 30:339-349.

80. Proctor WR, Poelchen W, Bowers BJ, Wehner JM, Messing RO, Dunwiddie TV, Ethanol differentially enhances hippocampal GABA A receptor-mediated responses in protein kinase C gamma (PKC gamma) and PKC epsilon null mice, Journal of Pharmacology and Experimental Therapeutics, 2003, 305:264-270.
81. Harris RA, McQuilkin SJ, Paylor R, Abeliovich A, Tonegawa S, Wehner JM, Mutant mice lacking the gamma isoform of protein kinase C show decreased behavioral actions of ethanol and altered function of gamma-aminobutyrate type A receptors, Proceedings of the National Academy of Sciences USA, 1995, 92:3658-3662.
82. Hodge CW, Mehmert KK, Kelley SP, McMahon T, Haywood A, Olive MF, Wang D, Sanchez-Perez AM, Messing RO, Supersensitivity to allosteric GABA(A) receptor modulators and alcohol in mice lacking PKCepsilon, Nature Neuroscience, 1999, 2:997-1002.
83. Hix DM, Bowers BJ, Miyamoto JH, Wehner JM, Open field activity and EtOH activation of gamma-PKC null mutants, Addiction Biology, 2003, 8:399-412.
84. Bowers BJ, Wehner JM, Ethanol consumption and behavioral impulsivity are increased in protein kinase C gamma null mutant mice, Journal of Neuroscience, 2001, 21:RC180.
85. Olive MF, McGeehan AJ, Kinder JR, McMahon T, Hodge CW, Janak PH, Messing RO, The mGluR5 antagonist 6-methyl-2-(phenylethynyl)pyridine decreases ethanol consumption via a protein kinase C epsilon-dependent mechanism, Molecular Pharmacology, 2005, 67:349-355.
86. Bowers BJ, McClure-Begley TD, Keller JJ, Paylor R, Collins AC, Wehner JM, Deletion of the alpha7 nicotinic receptor subunit gene results in increased sensitivity to several behavioral effects produced by alcohol, Alcoholism: Clinical and Experimental Research, 2005, 29:295-302.
87. Kralic JE, Criswell HE, Osterman JL, O'Buckley TK, Wilkie ME, Matthews DB, Hamre K, Breese GR, Homanics GE, Morrow AL, Genetic essential tremor in gamma-aminobutyric acidA receptor alpha1 subunit knockout mice, Journal of Clinical Investigation, 2005, 115:774-779.
88. Lallemand F, de Witte P, Ethanol induces higher BEC in CB1 cannabinoid receptor knockout mice while decreasing ethanol preference, Alcohol and Alcoholism, 2005, 40:54-62.
89. Blednov YA, Walker D, Osterndorf-Kahanek E, Harris RA, Mice lacking metabotropic glutamate receptor 4 do not show the motor stimulatory effect of ethanol, Alcohol, 2005, 34:251-259.

90. Boyce-Rustay JM, Holmes A, Functional roles of NMDA receptor NR2A and NR2B subunits in the acute intoxicating effects of ethanol in mice, Synapse, 2005, 56:222-225.
91. Tanaka K, Hashimoto H, Shintani N, Yamamoto A, Baba A, Reduced hypothermic and hypnotic responses to ethanol in PACAP-deficient mice, Regulatory Peptides, 2004, 123:95-98.
92. McKenzie-Quirk SD, Girasa KA, Allan AM, Miczek KA, 5-HT₃ receptors, alcohol and aggressive behavior in mice, Behavioural Pharmacology, 2005, 16:163-169.
93. Thanos PK, Rivera SN, Weaver K, Grandy DK, Rubinstein M, Umegaki H, Wang GJ, Hitzemann R, Volkow ND, Dopamine D₂R DNA transfer in dopamine D₂ receptor-deficient mice: Effects on ethanol drinking, Life Sciences, 2005, 77:130-139.
94. Newton PM, Orr CJ, Wallace MJ, Kim C, Shin HS, Messing RO, Deletion of N-type calcium channels alters ethanol reward and reduces ethanol consumption in mice, Journal of Neuroscience, 2004, 24:9862-9869.
95. Kovacs KM, Szakall I, O'Brien D, Wang R, Vinod KY, Saito M, Simonin F, Kieffer BL, Vadasz C, Decreased oral self-administration of alcohol in kappa-opioid receptor knock-out mice, Alcoholism: Clinical and Experimental Research, 2005, 29:730-8.
96. Pandey SC, Roy A, Zhang H, Xu T, Partial deletion of the cAMP response element-binding protein gene promotes alcohol-drinking behaviors, Journal of Neuroscience, 2004, 24:5022-5030.
97. Miyasaka K, Hosoya H, Takano S, Ohta M, Sekime A, Kanai S, Matsui T, Funakoshi A, Differences in ethanol ingestion between cholecystokinin-A receptor deficient and -B receptor deficient mice, Alcohol and Alcoholism, 2005, 40:176-180.
98. Maul B, Krause W, Pankow K, Becker M, Gembardt F, Alenina N, Walther T, Bader M, Siems WE, Central angiotensin II controls alcohol consumption via its AT₁ receptor, FASEB Journal, 2005, 19:1474-1481.
99. Thanos PK, Dimitrakakis ES, Rice O, Gifford A, Volkow ND, Ethanol self-administration and ethanol conditioned place preference are reduced in mice lacking cannabinoid CB₁ receptors, Behavioural Brain Research, 2005, 164:206-213.
100. Sharpe AL, Coste SC, Burkhart-Kasch S, Li N, Stenzel-Poore MP, Phillips TJ, Mice deficient in corticotropin-releasing factor receptor type 2 exhibit normal ethanol-associated behavior, Alcoholism: Clinical and Experimental Research, 2005, 29:1601-1609.

101. Blednov YA, Bergeson SE, Walker D, Ferreira VM, Kuziel WA, Harris RA, Perturbation of chemokine networks by gene deletion alters the reinforcing actions of ethanol, Behavioural Brain Research, 2005, 165:110-125.
102. Blednov YA, Walker D, Harris RA, Blockade of the leptin-sensitive pathway markedly reduces alcohol consumption in mice, Alcoholism: Clinical and Experimental Research, 2004, 28:1683-1692.
103. Ghozland S, Chu K, Kieffer BL, Roberts AJ, Lack of stimulant and anxiolytic-like effects of ethanol and accelerated development of ethanol dependence in mu-opioid receptor knockout mice, Neuropharmacology, 2005, 49:493-501.
104. Ivanova EA, Popova NK, Effect of monoamine oxidase A knockout on resistance to long-term exposure to ethanol, Bulletin of Experimental Biology and Medicine, 2002, 133:603-605.
105. Maas JW Jr, Vogt SK, Chan GC, Pineda VV, Storm DR, Muglia LJ, Calcium-stimulated adenylyl cyclases are critical modulators of neuronal ethanol sensitivity, Journal of Neuroscience, 2005, 25:4118-4126.
<NOTE: This one tested also the dowel test and vertical screen test, but there isn't room on the table for these entries>
106. McGough NN, He DY, Logrip ML, Jeanblanc J, Phamluong K, Luong K, Kharazia V, Janak PH, Ron D, RACK1 and brain-derived neurotrophic factor: a homeostatic pathway that regulates alcohol addiction, Journal of Neuroscience, 2004, 24:10542-10552.
107. Navarro M, Cubero I, Chen AS, Chen HY, Knapp DJ, Breese GR, Marsh DJ, Thiele TE, Effects of melanocortin receptor activation and blockade on ethanol intake: a possible role for the melanocortin-4 receptor, Alcoholism: Clinical and Experimental Research, 2005, 289:949-957.
108. Wand G, Levine M, Zweifel L, Schwindinger W, Abel T, The cAMP-protein kinase A signal transduction pathway modulates ethanol consumption and sedative effects of ethanol, Journal of Neuroscience, 2001, 21:5297-5303.
109. Stephens DN, Pistovcakova J, Worthing L, Atack JR, Dawson GR, Role of GABA(A) alpha5-containing receptors in ethanol reward: The effects of targeted gene deletion, and a selective inverse agonist, European Journal of Pharmacology, 2005, 526:240-250.
110. Sanchis-Segura C, Borchardt T, Vengeliene V, Zghoul T, Bachteler D, Gass P, Sprengel R, Spanagel R, Involvement of the AMPA receptor GluR-C subunit in alcohol-seeking behavior and relapse, Journal of Neuroscience, 2006, 26:1231-1238.

111. Vasiliou V, Ziegler TL, Bludeau P, Petersen DR, Gonzalez FJ, Deitrich RA, CYP2E1 and catalase influence ethanol sensitivity in the central nervous system, Pharmacogenetics and Genomics, 2006, 16:51-58.
112. Wallace MJ, Newton PM, Oyasu M, McMahon T, Chou WH, Connolly J, Messing RO, Acute functional tolerance to ethanol mediated by protein kinase C-epsilon, Neuropsychopharmacology, 2007, 32:127-136.
113. Boehm SL 2nd, Ponomarev I, Jennings AW, Whiting PJ, Rosahl TW, Garrett EM, Blednov YA, Harris RA, gamma-Aminobutyric acid A receptor subunit mutant mice: new perspectives on alcohol actions, Biochemical Pharmacology, 2004, 68:1581-1602.
114. Findlay GS, Wick MJ, Mascia MP, Wallace D, Miller GW, Harris RA, Blednov YA, Transgenic expression of a mutant glycine receptor decreases alcohol sensitivity of mice, Journal of Pharmacology and Experimental Therapeutics, 2002, 300:526-534.
115. Jeanblanc J, He DY, McGough NN, Logrip ML, Phamluong K, Janak PH, Ron D, The dopamine D3 receptor is part of a homeostatic pathway regulating ethanol consumption, Journal of Neuroscience, 2006, 26:1457-1464.
116. Yaka R, Tang KC, Camarini R, Janak PH, Ron D, Fyn kinase and NR2B-containing NMDA receptors regulate acute ethanol sensitivity but not ethanol intake or conditioned reward, Alcoholism: Clinical and Experimental Research, 2003, 27:1736-1742.
117. Basavarajappa BS, Yalamanchili R, Cravatt BF, Cooper TB, Hungund BL, Increased ethanol consumption and preference and decreased ethanol sensitivity in female FAAH knockout mice, Neuropharmacology, 2006, 50:834-844.
118. Fee JR, Knapp DJ, Sparta DR, Breese GR, Picker MJ, Thiele TE, Involvement of protein kinase A in ethanol-induced locomotor activity and sensitization, Neuroscience, 2006, 140:21-31.
119. Loney KD, Uddin RK, Singh SM, Analysis of metallothionein brain gene expression in relation to ethanol preference in mice using cosegregation and gene knockouts, Alcoholism: Clinical and Experimental Research, 2006, 30:15-25.
120. Olive MF, Mehmert KK, Koenig HN, Camarini R, Kim JA, Nannini MA, Ou CJ, Hodge CW, A role for corticotropin releasing factor (CRF) in ethanol consumption, sensitivity, and reward as revealed by CRF-deficient mice, Psychopharmacology, 2003, 165:181-187.
121. Cowen MS, Schroff KC, Gass P, Sprengel R, Spanagel R, Neurobehavioral effects of alcohol in AMPA receptor subunit (GluR1) deficient mice, Neuropharmacology, 2003, 45:325-333.

122. Ferraro FM 3rd, Sparta DR, Knapp DJ, Breese GR, Thiele TE, Increased consumption but not operant self-administration of ethanol in mice lacking the RIIbeta subunit of protein kinase A, Alcoholism: Clinical and Experimental Research, 2006, 30:825-835.
123. Primeaux SD, Wilson SP, Bray GA, York DA, Wilson MA, Overexpression of neuropeptide Y in the central nucleus of the amygdala decreases ethanol self-administration in "anxious" rats, Alcoholism: Clinical and Experimental Research, 2006, 30:791-801.
124. Werner C, Raivich G, Cowen M, Strelakova T, Sillaber I, Buters JT, Spanagel R, Hofmann F, Importance of NO/cGMP signalling via cGMP-dependent protein kinase II for controlling emotionality and neurobehavioural effects of alcohol, European Journal of Neuroscience, 2004, 20:3498-3506.
125. Isse T, Oyama T, Kitagawa K, Matsuno K, Matsumoto A, Yoshida A, Nakayama K, Nakayama K, Kawamoto T, Diminished alcohol preference in transgenic mice lacking aldehyde dehydrogenase activity, Pharmacogenetics, 2002, 12:621-626.
126. Sato Y, Seo N, Kobayashi E, Ethanol-induced hypnotic tolerance is absent in N-methyl-D-aspartate receptor epsilon-1 subunit knockout mice, Anesthesia and Analgesia, 2006, 103:117-120.
127. Besheer J, Lepoutre V, Mole B, Hodge CW, GABA(A) receptor regulation of voluntary ethanol drinking requires PKCepsilon, Synapse, 2006, 60:411-419.
128. Cai YQ, Cai GQ, Liu GX, Cai Q, Shi JH, Shi J, Ma SK, Sun X, Sheng ZJ, Mei ZT, Cui D, Guo L, Wang Z, Fei J, Mice with genetically altered GABA transporter subtype I (GAT1) expression show altered behavioral responses to ethanol, Journal of Neuroscience Research, 2006, 84:255-267.
129. Bonasera SJ, Chu HM, Brennan TJ, Tecott LH, A null mutation of the serotonin 6 receptor alters acute responses to ethanol, Neuropsychopharmacology, 2006, 31:1801-1813.
130. Boyce-Rustay JM, Holmes A, Ethanol-related behaviors in mice lacking the NMDA receptor NR2A subunit, Psychopharmacology, 2006, 187:455-466.
131. Metz AV, Chynoweth J, Allan AM, Influence of genetic background on alcohol drinking and behavioral phenotypes of 5-HT3 receptor over-expressing mice, Pharmacology Biochemistry and Behavior, 2006, 84:120-127.
132. Phillips TJ, Brown KJ, Burkhart-Kasch S, Wenger CD, Kelly MA, Rubinstein M, Grandy DK, Low MJ, Alcohol preference and sensitivity are markedly reduced in mice lacking dopamine D2 receptors, Nature Neuroscience, 1998, 1:610-615.

133. Crabbe JC, Wahlsten D, Dudek BC, Genetics of mouse behavior: interactions with laboratory environment, Science, 1999, 284:1670-1672.
134. Fee JR, Sparta DR, Knapp DJ, Breese GR, Picker MJ, Thiele TE, Predictors of high ethanol consumption in RIIbeta knock-out mice: assessment of anxiety and ethanol-induced sedation, Alcoholism: Clinical and Experimental Research, 2004, 28:1459-1468.
135. Szumlinski KK, Lominac KD, Oleson EB, Walker JK, Mason A, Dehoff MH, Klugmann M, Cagle S, Welt K, Doring M, Worley PF, Middaugh LD, Kalivas PW, Homer2 is necessary for EtOH-induced neuroplasticity, Journal of Neuroscience, 2005, 25:7054-7061 [erratum: 25(37):table of contents].
136. Offenhauser N, Castelletti D, Mapelli L, Soppo BE, Regondi MC, Rossi P, D'Angelo E, Frassoni C, Amadeo A, Tocchetti A, Pozzi B, Disanza A, Guarnieri D, Betsholtz C, Scita G, Heberlein U, Di Fiore PP, Increased ethanol resistance and consumption in eps8 knockout mice correlates with altered actin dynamics, Cell, 2006, 127:213-226.
137. Fernandez E, Koek W, Ran Q, Gerhardt GA, France CP, Strong R, Monoamine metabolism and behavioral responses to ethanol in mitochondrial aldehyde dehydrogenase knockout mice, Alcoholism: Clinical and Experimental Research, 2006, 30:1650-1658.
138. Werner DF, Blednov YA, Ariwodola OJ, Silberman Y, Logan E, Berry RB, Borghese CM, Matthews DB, Weiner JL, Harrison NL, Harris RA, Homanics GE, Knockin mice with ethanol-insensitive alpha1-containing gamma-aminobutyric acid type A receptors display selective alterations in behavioral responses to ethanol, Journal of Pharmacology and Experimental Therapeutics, 2006, 319:219-227.
139. Stork O, Kojima N, Stork S, Kume N, Obata K, Resistance to alcohol withdrawal-induced behaviour in Fyn transgenic mice and its reversal by ifenprodil, Molecular Brain Research, 2002, 105:126-135.
140. Cowen MS, Schumann G, Yagi T, Spanagel R, Role of Fyn tyrosine kinase in ethanol consumption by mice, Alcoholism: Clinical and Experimental Research, 2003, 27:1213-1219.
141. Abramov U, Raud S, Innos J, Koks S, Matsui T, Vasar E, Gender specific effects of ethanol in mice, lacking CCK(2) receptors, Behavioural Brain Research, 2006, 175:149-156.
142. Boyce-Rustay JM, Wiedholz LM, Millstein RA, Carroll J, Murphy DL, Daws LC, Holmes A, Ethanol-related behaviors in serotonin transporter knockout mice, Alcoholism: Clinical and Experimental Research, 2006, 30:1957-1965.

143. Caldwell HK, Stewart J, Wiedholz LM, Millstein RA, Iacangelo A, Holmes A, Young WS 3rd, Wersinger SR, The acute intoxicating effects of ethanol are not dependent on the vasopressin 1a or 1b receptors, Neuropeptides, 2006, 40:325-337.
144. Zghoul T, Abarca C, Sanchis C, Albrecht U, Schumann G, Spanagel R, Ethanol self-administration and reinstatement of ethanol-seeking behavior in *Per1^{Brdm1}* mutant mice, Psychopharmacology, 2007, 190:13-19.
145. June HL Sr, Foster KL, Eiler WJA 2nd, Goergen J, Cook JB, Johnson N, Mensah-Zoe B, Simmons JO, June HL Jr, Yin W, Cook JM, Homanics GE, Dopamine and benzodiazepine-dependent mechanisms regulate the EtOH-enhanced locomotor stimulation in the GABA-A alpha-1 subunit null mutant mice, Neuropsychopharmacology, 2007, 32:137-152.
146. Faber F, Gembardt F, Sun X, Mizutani S, Siems WE, Walther T, Lack of angiotensin II conversion to angiotensin III increases water but not alcohol consumption in aminopeptidase A-deficient mice, Regulatory Peptides, 2006, 136:130-137.
147. Savelieva KV, Caudle WM, Miller GW, Altered ethanol-associated behaviors in vesicular monoamine transporter heterozygote knockout mice, Alcohol, 2006, 40:87-94.
148. Blednov YA, Walker D, Martinez M, Harris RA, Reduced alcohol consumption in mice lacking preprodynorphin, Alcohol, 2006, 40:73-86.
149. Newton PM, Messing RO, Increased sensitivity to the aversive effects of ethanol in PKC ϵ null mice revealed by place conditioning, Behavioral Neuroscience, 2007, 121:439-442.
150. Tornberg J, Segerstrale M, Kuleskaya N, Voikar V, Taira T, Airaksinen MS, KCC2-deficient mice show reduced sensitivity to diazepam, but normal alcohol-induced motor impairment, gaboxadol-induced sedation, and neurosteroid-induced hypnosis, Neuropsychopharmacology, 2007, 32:911-918.
151. Jeans AF, Oliver PL, Johnson R, Capogna M, Vikman J, Molnar Z, Babbs A, Partridge CJ, Salehi A, Bengtsson M, Eliasson L, Rorsman P, Davies KE, A dominant mutation in Snap25 causes impaired vesicle trafficking, sensorimotor gating, and ataxia in the blind-drunk mouse, Proceedings of the National Academy of Sciences USA, 2007, 104:2431-2436.
152. Moore R, Krstew EV, Kirchhoff J, Davisson RL, Lawrence AJ, Central overexpression of angiotensin AT(1A) receptors prevents dopamine D(2) receptor regulation of alcohol consumption in mice, Alcoholism: Clinical and Experimental Research, 2007, 31:1128-1137.
153. Sanchis-Segura C, Cline B, Jurd R, Rudolph U, Spanagel R, Etomidate and propofol-hyposensitive GABAA receptor beta3(N265M) mice show little changes in acute alcohol

- sensitivity but enhanced tolerance and withdrawal, Neuroscience Letters, 2007, 416:275-278.
154. Warnault V, Houchi H, Barbier E, Pierrefiche O, Vilpoux C, Ledent C, Daoust M, Naassila M, The lack of CB1 receptors prevents neuroadaptations of both NMDA and GABA-A receptors after chronic ethanol exposure, Journal of Neurochemistry, 2007, 102:741-752.
 155. Chu K, Koob GF, Cole M, Zorrilla EP, Roberts AJ, Dependence-induced increases in ethanol self-administration in mice are blocked by the CRF1 receptor antagonist antalarmin and by CRF1 receptor knockout, Pharmacology Biochemistry and Behavior, 2007, 86:813-821.
 156. Blednov YA, Cravatt BF, Boehm SL 2nd, Walker D, Harris RA, Role of endocannabinoids in alcohol consumption and intoxication: studies of mice lacking fatty acid amide hydrolase, Neuropsychopharmacology, 2007, 32:1570-1582.
 157. Duncan EA, Sorrell JE, Adamantidis A, Rider T, Jandacek RJ, Seeley RJ, Lakaye B, Woods SC, Alcohol drinking in MCH receptor-1-deficient mice, Alcoholism: Clinical and Experimental Research, 2007, 31:1325-1337.
 158. Sparta DR, Fee JR, Knapp DJ, Breese GR, Thiele TE, Elevated anxiety-like behavior following ethanol exposure in mutant mice lacking neuropeptide Y (NPY), Drug and Alcohol Dependence, 2007, 90:297-300.
 159. Vinod KY, Sanguino E, Yalamanchili R, Manzanares J, Hungund BL, Manipulation of fatty acid amid hydrolase functional activity alters sensitivity and dependence to ethanol, Journal of Neurochemistry, 2008, 104:233-243.
 160. Chandra D, Werner DF, Liang J, Suryanarayanan A, Harrison NL, Spigelman I, Olsen RW, Homanics GE, Normal acute behavioral responses to moderate/high dose ethanol in GABA-A receptor $\alpha 4$ subunit knockout mice, Alcoholism: Clinical and Experimental Research, 2008, 32:10-18.
 161. Sakoori K, Murphy NP, Endogenous nociceptin (orphanin FQ) suppresses basal hedonic state and acute reward responses to methamphetamine and ethanol, but facilitates chronic responses, Neuropsychopharmacology, 2008, 33:877-891.
 162. Blednov YA, Walker D, Martinez M, Levine M, Damak S, Margolskee RF, Perception of sweet taste is important for voluntary alcohol consumption in mice, Genes Brain and Behavior, 2008, 7:1-13.
 163. Stemmelin J, Cohen C, Terranova JP, Lopez-Grancha M, Pichat P, Bergis O, Decobert M, Santucci V, Françon D, Alonso R, Stahl SM, Keane P, Avenet P, Scatton B, le Fur G,

- Griebel G. Stimulation of the beta3-Adrenoceptor as a novel treatment strategy for anxiety and depressive disorders. Neuropsychopharmacology, 2008, 33:574-587.
164. Szumlinski KK, Ary AW, Lominac KD, Klugmann M, Kippin TE. Accumbens Homer2 overexpression facilitates alcohol-induced neuroplasticity in C57BL/6J mice. Neuropsychopharmacology, 2008, 33:1365-1378.
165. Vinod KY, Yalamanchili R, Thanos PK, Vadasz C, Cooper TB, Volkow ND, Hungund BL, Genetic and pharmacological manipulations of the CB1 receptor alter ethanol preference and dependence in ethanol preferring and nonpreferring mice, Synapse, 2008, 62:574-581.
166. Sanbe A, Takagi N, Fujiwara Y, Yamauchi J, Endo T, Mizutani R, Takeo S, Tsujimoto G, Tanoue A. Alcohol preference in mice lacking the Avpr1a vasopressin receptor. American Journal of Physiology Regulatory Integrative and Comparative Physiology, 2008, 294:R1482-R1490.
167. Pastor R, McKinnon CS, Scibelli AC, Burkhart-Kasch S, Reed C, Ryabinin AE, Coste SC, Stenzel-Poore MP, Phillips TJ. Corticotropin-releasing factor-1 receptor involvement in behavioral neuroadaptation to ethanol: a urocortin-1-independent mechanism. Proceedings of the National Academy of Sciences USA, 2008, 105:9070-9075.
168. Björk K, Rimondini R, Hansson AC, Terasmaa A, Hyttiä P, Heilig M, Sommer WH. Modulation of voluntary ethanol consumption by beta-arrestin 2. FASEB Journal, 2008, 22:2552-2560.
169. Grisel JE, Bartels JL, Allen SA, Turgeon VL. Influence of β -endorphin on anxious behavior in mice: interaction with EtOH. Psychopharmacology, 2008, 200:105-115.
170. Palachik B, Chen YC, Enoch AJ, Karlsson RM, Mishina M, Holmes A. Role of major NMDA or AMPA receptor subunits in MK-801 potentiation of ethanol intoxication. Alcoholism: Clinical and Experimental Research, 2008, 32:1479-1492.
171. Itzhak Y, Anderson KL. Ethanol-induced behavioral sensitization in adolescent and adult mice: role of the nNOS gene. Alcoholism: Clinical and Experimental Research, 2008, 32:1839-1848.
172. Choi DS, Wei W, Deitchman JK, Kharazia VN, Lesscher HMB, McMahon T, Wang D, Qi ZH, Sieghart W, Zhang C, Shokat KM, Mody I, Messing RO. Protein kinase C δ regulates ethanol intoxication and enhancement of GABA-stimulated tonic current. Journal of Neuroscience, 2008, 28:11890-11899.
173. Racz I, Schurmann B, Karpushova A, Reuter M, Cichon S, Montag C, Furst R, Schutz C, Franke PE, Strohmaier J, Wienker TF, Terenius L, Osby U, Gunnar A, Maier W, Bilkei-

- Gorzo A, Nothen M, Zimmer A. The opioid peptides enkephalin and beta-endorphin in alcohol dependence. Biological Psychiatry, 2008, 64:989-997.
174. Houchi H, Warnault V, Barbier E, Dubois C, Pierrefiche O, Ledent C, Daoust M, Naassila M. Involvement of A2A receptors in anxiolytic, locomotor and motivational properties of ethanol in mice. Genes, Brain and Behavior, 2008, 7:887-898.
175. Blednov YA, Harris RA. Metabotropic glutamate receptor 5 (mGluR5) regulation of ethanol sedation, dependence and consumption: relationship to acamprosate actions. International Journal of Neuropsychopharmacology, 2008, 11:775-793.
176. Bird MK, Kirchhoff J, Djouma E, Lawrence AJ. Metabotropic glutamate 5 receptors regulate sensitivity to ethanol in mice. International Journal of Neuropsychopharmacology, 2008, 11:765-774.
177. Tammimaki A, Forsberg MM, Karayiorgou M, Gogos JA, Mannisto PT. Increase in free choice oral ethanol self-administration in catechol-*O*-methyltransferase gene-disrupted male mice. Basic and Clinical Pharmacology and Toxicology, 2008, 103:297-304.
178. Ellingson JM, Silbaugh BC, Brassler SM. Reduced oral ethanol avoidance in mice lacking transient receptor potential channel vanilloid receptor 1. Behavior Genetics, 2009, 39:62-72.
179. Martin GE, Hendrickson LM, Penta KL, Friesen RM, Pietrzykowski AZ, Tapper AR, Treistman SN. Identification of a BK channel auxiliary protein controlling molecular and behavioral tolerance to alcohol. Proceedings of the National Academy of Sciences USA, 2008, 105:17543-17548.
180. Newton PM, Zeng L, Wang V, Connolly J, Wallace MJ, Kim C, Shin HS, Belardetti F, Snutch TP, Messing RO. A blocker of N- and T-type voltage-gated calcium channels attenuates ethanol-induced intoxication, place preference, self-administration, and reinstatement. Journal of Neuroscience, 2008, 28:11712-11719.
181. Kapfhamer D, Bettinger JC, Davies AG, Eastman CL, Smail EA, Heberlein U, McIntire SL. Loss of RAB-3/A in *Caenorhabditis elegans* and the mouse affects behavioral response to ethanol. Genes, Brain and Behavior, 2008, 7:669-676.
182. Olmstead MC, Ouagazzal AM, Kieffer BL. Mu and delta opioid receptors oppositely regulate motor impulsivity in the signaled nose poke task. PLOS One, 2009, 4:e4410.
183. Blednov YA, Harris RA. Deletion of vanilloid receptor (TRPV1) in mice alters behavioral effects of ethanol. Neuropharmacology, 2009, 56:814-820.

184. Berry RB, Chandra D, Diaz-Granados JL, Homanics GE, Matthews DB. Investigation of ethanol-induced impairment of spatial memory in γ 2 heterozygous knockout mice. Neuroscience Letters, 2009, 455:84-87.
185. Correa M, Viaggi C, Escrig MA, Pascual M, Guerri C, Vaglini F, Aragon CM, Corsini GU. Ethanol intake and ethanol-induced locomotion and locomotor sensitization in Cyp2e1 knockout mice. Pharmacogenetics and Genomics, 2009, 19:217-225.
186. Harrison SJ, Nobrega JN. Differential susceptibility to ethanol and amphetamine sensitization in dopamine D3 receptor-deficient mice. Psychopharmacology, 2009, 204:49-59.
187. Ting-A-Kee R, Dockstader C, Heinmiller A, Grieder T, van der Kooy D. GABA-A receptors mediate the opposing roles of dopamine and the tegmental pedunculopontine nucleus in the motivational effects of ethanol. European Journal of Neuroscience, 2009, 29:1235-1244.
188. Navarro M, Cubero I, Ko L, Thiele TE. Deletion of agouti-related protein blunts ethanol self-administration and binge-like drinking in mice. Genes Brain and Behavior, 2009, 8:450-458.
189. Jerlhag E, Egecioglu E, Landgren S, Salomé N, Heilig M, Moechars D, Datta R, Perrissoud D, Dickson SL, Engel JA. Requirement of central ghrelin signaling for alcohol reward. Proceedings of the National Academy of Sciences USA, 2009, 106:11318-11323.
190. Itzhak Y, Roger-Sánchez C, Anderson KL. Role of the nNOS gene in ethanol-induced conditioned place preference in mice. Alcohol, 2009, 43:285-291.
191. Carnicella S, Ahmadiantehrani S, He DY, Nielsen CK, Bartlett SE, Janak PH, Ron D. Cabergoline decreases alcohol drinking and seeking behaviors via glial cell line-derived neurotrophic factor. Biological Psychiatry, 2009, 66:146-153.
192. Kozell LB, Walter NA, Milner LC, Wickman K, Buck KJ. Mapping a barbiturate withdrawal locus to a 0.44 Mb interval and analysis of a novel null mutant identify a role for Kcnj9 (GIRK3) in withdrawal from pentobarbital, zolpidem, and ethanol. Journal of Neuroscience, 2009, 29:11662-11673.
193. van Rijn RM, Whistler JL. The delta-1 opioid receptor is a heterodimer that opposes the actions of the delta-2 receptor on alcohol intake. Biological Psychiatry, 2009, 66:777-784.
194. Lesscher HMB, Wallace MJ, Zeng L, Wang V, Deitchman JK, McMahon T, Messing RO, Newton PM. Amygdala protein kinase C epsilon controls alcohol consumption. Genes, Brain and Behavior, 2009, 8:493-499.

195. Gulick D, Gould TJ. The hippocampus and cingulate cortex differentially mediate the effects of nicotine on learning versus on ethanol-induced learning deficits through different effects at nicotinic receptors. Neuropsychopharmacology, 2009, 34:2167-2179.
196. Blednov YA, Walker DL, Iyer SV, Homanics G, Harris AR [sic]. mice lacking Gad2 show altered behavioral effects of ethanol, flurazepam and gabaxadol. Addiction Biology, 2010, 15:45-61.
197. Karatayev O, Baylan J, Leibowitz SF. Increased intake of ethanol and dietary fat in galanin overexpressing mice. Alcohol, 2009, 43:571-580.
198. Karatayev O, Baylan J, Weed V, Chang S, Wynick D, Leibowitz SF. Galanin knockout mice show disturbances in ethanol consumption and expression of hypothalamic peptides that stimulate ethanol intake. Alcoholism: Clinical and Experimental Research, 2010, 34:72-80.
199. Morice E, Denis C, Giros B, Nosten-Bertrand M. Evidence of long-term expression of behavioral sensitization to both cocaine and ethanol in dopamine transporter knockout mice. Psychopharmacology, 2010, 208:57-66.
200. Nuutinen S, Karlstedt K, Aitta-Aho T, Korpi ER, Panula P. Histamine and H3 receptor-dependent mechanisms regulate ethanol stimulation and conditioned place preference in mice. Psychopharmacology, 2010, 208:75-86.
201. Mutschler J, Bilbao A, von der Goltz C, Demiralay C, Jahn H, Wiedemann K, Spanagel R, Kiefer F. Augmented stress-induced alcohol drinking and withdrawal in mice lacking functional natriuretic peptide-A receptors. Alcohol and Alcoholism, 2010, 45:13-16.
202. Tanaka K, Kunishige-Yamamoto A, Hashimoto H, Shintani N, Hayata A, Baba A. Increased ethanol preference and serotonin 1A receptor-dependent attenuation of ethanol-induced hypothermia in PACAP-deficient mice. Biochemical and Biophysical Research Communications, 2010, 391:773-777.
203. Chen J, Nam HW, Lee MR, Hinton DJ, Choi S, Kim T, Kawamura T, Janak PH, Choi DS. Altered glutamatergic neurotransmission in the striatum regulates ethanol sensitivity and intake in mice lacking ENT1. Behavioural Brain Research, 2010, 208:636-642.
204. Lee MR, Hinton DJ, Song JY, Lee KW, Choo C, Johng H, Unal SS, Richelson E, Choi DS. Neurotensin receptor type 1 regulates ethanol intoxication and consumption in mice. Pharmacology Biochemistry and Behavior, 2010, 95:235-241.
205. Thorsell A, Schank JR, Singley E, Hunt SP, Heilig M. Neurokinin-1 receptors (NK1R:s), alcohol consumption, and alcohol reward in mice. Psychopharmacology, 2010, 209:103-111.

206. Sharpe AL, Low MJ. Proopiomelanocortin peptides are not essential for development of ethanol-induced behavioral sensitization. Alcoholism: Clinical and Experimental Research, 2009, 33:1202-1207.
207. Kamens HM, McKinnon CS, Li N, Helms ML, Belknap JK, Phillips TJ. The $\alpha 3$ subunit gene of the nicotinic acetylcholine receptor is a candidate gene for ethanol stimulation. Genes Brain and Behavior, 2009, 8:600-609.
208. Sperling RE, Gomes SM, Sypek EI, Carey AN, McLaughlin JP. Endogenous kappa-opioid mediation of stress-induced potentiation of ethanol-conditioned place preference and self-administration. Psychopharmacology, 2010, 210:199-209.
209. Hendrickson LM, Zhao-Shea R, Pang X, Gardner PD, Tapper AR. Activation of $\alpha 4^*$ nAChRs is necessary and sufficient for varenicline-induced reduction of alcohol consumption. Journal of Neuroscience, 2010, 30:10169-10176.
210. Repunte-Canonigo V, Berton F, Cottone P, Reifel-Miller A, Roberts AJ, Morales M, Francesconi W, Sanna PP. A potential role for adiponectin receptor 2 (AdipoR2) in the regulation of alcohol intake. Brain Research, 2010a, 1339:11-17.
211. Mathews TA, Brookshire BR, Budygin EA, Hamre K, Goldowitz D, Jones SR. Ethanol-induced hypoactivity is associated with hypodopaminergia in the 22-TNJ ENU-mutated mouse. Alcohol, 2009, 43:421-431.
212. Speca DJ, Chihara D, Ashique AM, Bowers MS, Pierce-Shimomura JT, Lee J, Rabbee N, Speed TP, Gularte RJ, Chitwood J, Medrano JF, Liao M, Sonner JM, Eger EI 2nd, Peterson AS, McIntire SL. Conserved role of *unc-79* in ethanol responses in *Lightweight* mutant mice. PLoS Genetics, 2010, 6:e1001057.
213. Repunte-Canonigo V, van der Stap LD, Chen J, Sabino V, Wagner U, Zorrilla EP, Schumann G, Roberts AJ, Sanna PP. Genome-wide gene expression analysis identifies K-ras as a regulator of alcohol intake. Brain Research, 2010b, 1339:1-10.
214. Nam HW, Lee MR, Hinton DJ, Choi DS. Reduced effect of NMDA glutamate receptor antagonist on ethanol-induced ataxia and striatal glutamate levels in mice lacking ENT1. Neuroscience Letters, 2010, 479:277-281.
215. Downing C, Marks MJ, Larson C, Johnson TE. The metabotropic glutamate receptor subtype 5 mediates sensitivity to the sedative properties of ethanol. Pharmacogenetics and Genomics, 2010, 20:553-564.
216. Brassler SM, Norman MB, Lemon CH. T1r3 taste receptor involvement in gustatory neural responses to ethanol and oral ethanol preference. Physiological Genomics, 2010, 41:232-243.

217. Barfield ET, Barry SM, Hodgins HB, Thompson BM, Allen SS, Grisel JE. β -endorphin mediates behavioral despair and the effect of ethanol on the tail suspension test in mice. Alcoholism: Clinical and Experimental Research, 2010, 34:1066-1072.
218. Martin LJ, Zurek AA, Bonin RP, Oh GHT, Kim JH, Mount HTJ, Orser BA. The sedative but not the memory-blocking properties of ethanol are modulated by α 5-subunit-containing γ -aminobutyric acid type A receptors. Behavioural Brain Research, 2011, 217:379-385.
219. Kim JH, Karpyak VM, Biernacka JM, Nam HW, Lee MR, Preuss UW, Zill P, Yoon G, Colby C, Mrazek DA, Choi DS. Functional role of the polymorphic 647 T/C variant of ENT1 (*SLC29A1*) and its association with alcohol withdrawal seizures. PLoS One, 2011, 6:e16331.
220. Gibb SL, Jeanblanc J, Barak S, Yowell QV, Yaka R, Ron D. Lyn kinase regulates mesolimbic dopamine release: implication for alcohol reward. Journal of Neuroscience, 2011, 31:2180-2187.
221. Toth ME, Gonda S, Vigh L, Santha M. Neuroprotective effect of small heat shock protein, Hsp27, after acute and chronic alcohol administration. Cell Stress and Chaperones, 2010, 15:807-817.
222. Lee MR, Hinton DJ, Unal SS, Richelson E, Choi DS. Increased ethanol consumption and preference in mice lacking neurotensin receptor type 2. Alcoholism: Clinical and Experimental Research, 2011, 35:99-107.
223. Linden AM, Schmitt U, Leppa E, Wulff P, Wisden W, Luddens H, Korpi ER. Ro 15-4513 antagonizes alcohol-induced sedation in mice through $\alpha\beta\gamma$ 2-type GABA_A receptors. Frontiers in Neuroscience, 2011, 5:3.
224. Navarro M, Lerma-Cabrera JM, Cravajal F, Lowery EG, Cubero I, Thiele TE. Assessment of voluntary ethanol consumption and the effects of a melanocortin (MC) receptor agonist on ethanol intake in mutant C57BL/6J mice lacking the MC-4 receptor. Alcoholism: Clinical and Experimental Research, 2011, 35:1058-1066.
225. Nam HW, Lee MR, Zhu Y, Wu J, Hinton DJ, Choi S, Kim T, Hammack N, Yin JCP, Choi DS. Type 1 equilibrative nucleoside transporter regulates ethanol drinking through accumbal N-methyl-D-aspartate receptor signaling. Biological Psychiatry, 2011, 69:1043-1051.
226. Mittleman G, Call SB, Cockroft JL, Goldowitz D, Matthews DB, Blaha CD. Dopamine dynamics associated with, and resulting from, schedule-induced alcohol self-administration: analyses in dopamine transporter knockout mice. Alcohol, 2011, 45:325-339.

227. Kost S, Sun C, Xiong W, Graham K, Cass CE, Young JD, Albeni BC, Parkinson FE. Behavioral effects of elevated expression of human equilibrative nucleoside transporter 1 in mice. Behavioural Brain Research, 2011, 224:44-49.
228. Kim KS, Kim H, Baek IS, Lee KW, Han PL. Mice lacking adenylyl cyclase type 5 (AC5) show increased ethanol consumption and reduced ethanol sensitivity. Psychopharmacology, 2011, 215:391-398.
229. Kamens HM, Andersen J, Picciotto MR. Modulation of ethanol consumption by genetic and pharmacological manipulation of nicotinic acetylcholine receptors in mice. Psychopharmacology, 2010, 208:613-626.
230. Jerlhag E, Landgren S, Egecioglu E, Dickson SL, Engel JA. The alcohol-induced locomotor stimulation and accumbal dopamine release is suppressed in ghrelin knockout mice. Alcohol, 2011, 45:341-347.
231. Steffensen SC, Bradley KD, Hansen DM, Wilcox JD, Wilcox RS, Allison DW, Merrill CB, Edwards JG. The role of connexin-36 gap junctions in alcohol intoxication and consumption. Synapse, 2011, 65:695-707.
232. Desrivieres S, Pronko SP, Lourdasamy A, Ducci F, Hoffman PL, Wodarz N, Ridinger M, Rietschel M, Zelenika D, Lathrop M, Schumann G, Tabakoff B. Sex-specific role for adenylyl cyclase type 7 in alcohol dependence. Biological Psychiatry, 2011, 69:1100-1108.
233. Mittleman G, Call SB, Cockroft JL, Goldowitz D, Matthews DB, Blaha CD. Dopamine dynamics associated with, and resulting from, schedule-induced alcohol self-administration: analyses in dopamine transporter knockout mice. Alcohol, 2011, 45:325-339.
234. Camp MC, Feyder M, Ihne J, Palachick B, Hurd B, Kaerlsson RM, Noronha B, Chen YC, Coba MP, Grant SGN, Holmes A. A novel role for PSD-95 in mediating ethanol intoxication, drinking and place preference. Addiction Biology, 2011, 16:428-439.
235. Harris RA, Osterndorff-Kahanek E, Ponomarev I, Homanics GE, Blednov YA. Testing the silence of mutations: transcriptomic and behavioral studies of GABAA receptor $\alpha 1$ and $\alpha 2$ subunit knock-in mice. Neuroscience Letters, 2011, 488:31-35.
236. Hicklin TR, Wu PH, Radcliffe RA, Freund RK, Goebel-Goody SM, Correa PR, Proctor WR, Lombroso PJ, Browning MD. Alcohol inhibition of the NMDA receptor function, long-term potentiation, and fear learning requires striatal-enriched protein tyrosine phosphatase. Proceedings of the National Academy of Sciences USA, 2011, 108:6650-6655.

237. Brager AJ, Prosser RA, Glass JD. Circadian and acamprosate modulation of elevated ethanol drinking in mPer2 clock gene mutant mice. Chronobiology International, 2011, 28:664-672.
238. Bulwa ZB, Sharlin JA, Clark PJ, Bhattacharya TK, Kilby CN, Wang Y, Rhodes JS. Increased consumption of ethanol and sugar water in mice lacking the dopamine D2 long receptor. Alcohol, 2011, 45:631-639.
239. Iyer SV, Benavides RA, Chandra D, Cook JM, Rallapalli S, June HL, Homanics GE. α 4-Containing GABA_A receptors are required for antagonism of ethanol-induced motor incoordination and hypnosis by the imidazobenzodiazepine Ro15-4513. Frontiers in Pharmacology, 2011, 2:18.
240. Lasek AW, Lim J, Kliethermes CL, Berger KH, Joslyn G, Brush G, Xue L, Robertson M, Moore MS, Vranizan K, Morris SW, Schuckit MA, White RL, Heberlein U. An evolutionary conserved role for anaplastic lymphoma kinase in behavioral responses to ethanol. PLoS One, 2011, 6:e22636.
241. Dong L, Bilbao A, Laucht M, Henriksson R, Yakovleva T, Ridinger M, Desrivieres S, Clarke TK, Lourdasamy A, Smolka MN, Cichon S, Blomeyer D, Treutlein J, Peralta-Lenz S, Witt S, Leonardi-Essmann F, Wodarz N, Ill P, Soyka M, Albrecht U, Rietschel M, Lathrop M, Bakalkin G, Spanagel R, Schumann G. Effects of the circadian rhythm gene Period 1 (Per1) on psychosocial stress-induced alcohol drinking. American Journal of Psychiatry, 2011, 168:1090-1098.
242. Badanich KA, Doremus-Fitzwater TL, Mulholland PJ, Randall PK, Delpire E, Becker HC. NR2B-deficient mice are more sensitive to the locomotor stimulant and depressant effects of ethanol. Genes, Brain and Behavior, 2011, 10:805-816.
243. Blednov YA, Ponomarev I, Geil C, Bergeson S, Koob GF, Harris RA. Neuroimmune regulation of alcohol consumption: behavioral validation of genes obtained from genomic studies. Addiction Biology, 2011, 17:108-120.
244. Brager A, Prosser RA, Glass JD. Acamprosate-responsive brain sites for suppression of ethanol intake and preference. American Journal of Physiology: Regulatory, Integrative and Comparative Physiology, 2011, 301:R1032-R1043.
245. Du X, Elberger AJ, Matthews DB, Hamre KM. Heterozygous deletion of NR1 subunit of the NMDA receptor alters ethanol-related behaviors and regional expression of NR2 subunits in the brain. Neurotoxicology and Teratology, 2012, 34:177-186.
246. Jaholkowski P, Mierzejewski P, Zatorski P, Scinska A, Sienkiewicz-Jarosz H, Kaczmarek L, Samochowicz J, Filipkowski RK, Bienkowski P. Increased ethanol intake and preference in cyclin D2 knockout mice. Genes Brain and Behavior, 2011, 10:551-556.

247. Kaur S, Li J, Stenzel-Poore MP, Ryabinin AE. Corticotropin-releasing factor acting on corticotropin-releasing factor receptor type 1 is critical for binge alcohol drinking in mice. Alcoholism: Clinical and Experimental Research, 2012, 36:369-376.
248. Giardino WJ, Cocking DL, Kaur S, Cunningham CL, Ryabinin AE. Urocortin-1 within the centrally-projecting Edinger-Westphal nucleus is critical for ethanol preference. PLoS One, 2011, 6:e26997.
249. Lee MR, Hinton DJ, Wu J, Mishra PK, Port JD, Macura SI, Choi DS. Acamprosate reduces ethanol drinking behaviors and alters the metabolite profile in mice lacking ENT1. Neuroscience Letters, 2011, 490:90-95.
250. Molander A, Vengeliene V, Heilig M, Wurst W, Deussing JM, Spaangel R. Brain-specific inactivation of the Crhr1 gene inhibits post-dependent and stress-induced alcohol intake, but does not affect relapse-like drinking. Neuropsychopharmacology, 2012, 37:1047-1056.
251. Nguyen K, Tseng A, Marquez P, Hamid A, Lutfy K. The role of endogenous dynorphin in ethanol-induced state-dependent CPP. Behavioural Brain Research, 2012, 227:58-63.
252. Nuutinen S, Lintunen M, Vanhanen J, Ojala T, Rozov S, Panula P. Evidence for the role of histamine H3 receptor in alcohol consumption and alcohol reward in mice. Neuropsychopharmacology, 2011, 36:2030-2040.
253. Pastor R, Reed C, Burkhardt-Kasch S, Li N, Sharpe AL, Coste SC, Stenzel-Poore MP, Phillips TJ. Ethanol concentration-dependent effects and the role of stress on ethanol drinking in corticotropin-releasing factor type 1 and double type 1 and 2 receptor knockout mice. Psychopharmacology, 2011, 218:169-177.
254. Wu Y, Lousberg EL, Moldenhauer LM, Hayball JD, Coller JK, Rice KC, Watkins LR, Somogyi AA, Hutchinson MR. Inhibiting the TLR4-MyD88 signalling cascade by genetic or pharmacological strategies reduces acute alcohol-induced sedation and motor impairment in mice. British Journal of Pharmacology, 2012, 165:1319-1329.
255. Cushman JD, Moore MD, Jacobs NS, Olsen RW, Fanselow MS. Behavioral pharmacogenetic analysis on the role of the $\alpha 4$ GABA_A receptor subunit in the ethanol-mediated impairment of hippocampus-dependent contextual learning. Alcoholism: Clinical and Experimental Research, 2011, 35:1948-1959.
256. Femenia T, Manzanares J. Increased ethanol intake in prodynorphin knockout mice is associated to changes in opioid receptor function and dopamine transmission. Addiction Biology, 2011, 17:322-337.

257. Hayes DM, Fee JR, McCown TJ, Knapp DJ, Breese GR, Cubero I, Carvajal F, Lerma-Cabrera JM, Navarro M, Thiele TE. Neuropeptide Y signaling modulates the expression of ethanol-induced behavioral sensitization in mice. Addiction Biology, 2011, 17:338-350.
258. Bhandari P, Hill JS, Farris SP, Costin B, Martin I, Chan CL, Alaimo JT, Bettinger JC, Davies AG, Miles MF, Grotewiel M. Chloride intracellular channels modulate acute ethanol behaviors in *Drosophila*, *Caenorhabditis elegans* and mice. Genes, Brain and Behavior, 2012, 11:387-397.
259. Petit-Paitel A, Menard B, Guyon A, Beringue V, Nahon JL, Zsurger N, Chabry J. Prior protein is a key determinant of alcohol sensitivity through the modulation of N-methyl-D-aspartate receptor (NMDAR) activity. PLoS One, 2012, 7:e34691.
260. Pastor R, Reed C, Meyer PJ, McKinnon C, Ryabinin AE, Phillips TJ. Role of corticotropin-releasing factor and corticosterone in behavioral sensitization to ethanol. Journal of Pharmacology and Experimental Therapeutics, 2012, 341:455-463.
261. Karlsson RM, Adermark L, Molander A, Perreau-Lenz S, Singley E, Solomon M, Holmes A, Tanaka K, Lovinger DM, Spanagel R, Heilig M. Reduced alcohol intake and reward associated with impaired endocannabinoid signaling in mice with a deletion of the glutamate transporter GLAST. Neuropharmacology, 2012, 63:181-189.
262. Gallego X, Ruiz-Medina J, Valverde O, Molas S, Robles N, Sabrià J, Crabbe JC, Dierssen M. Transgenic over expression of nicotinic receptor alpha 5, alpha 3, and beta 4 subunit genes reduces ethanol intake in mice. Alcohol, 2012, 46:205-215.
263. Kamens HM, Hoft NR, Cox RJ, Miyamoto JH, Ehringer MA. The $\alpha 6$ nicotinic acetylcholine receptor subunit influences ethanol-induced sedation. Alcohol, 2012, 46:463-471.
264. Blednov YA, Benavidez JM, Black M, Chandra D, Homanics GE, Rudolph U, Harris RA. Linking GABA_A receptor subunits to alcohol-induced conditioned taste aversion and recovery from acute alcohol intoxication. Neuropharmacology, 2012, 67C:46-56.
265. Ary AW, Cozzoli DK, Finn DA, Crabbe JC, Dehoff MH, Worley PF, Szumlinski KK. Ethanol up-regulates nucleus accumbens neuronal activity dependent pentraxin (Narp): implications for alcohol-induced behavioral plasticity. Alcohol, 2012, 46:377-387.
266. Dixon CI, Walker SE, King SL, Stephens DN. Deletion of the *gabra2* gene results in hypersensitivity to the acute effects of ethanol but does not alter ethanol self administration. PLoS One, 2012,7:e47135.

267. Kim KS, Kim H, Park SK, Han PL. The dorsal striatum expressing adenylyl cyclase-5 controls behavioral sensitivity of the righting reflex to high-dose ethanol. Brain Research, 2012, 1489:27-36.
268. Maul B, Becker M, Gembardt F, Becker A, Schultheiss HP, Siems WE, Walther T. Genetic deficiency in neprilysin or its pharmacological inhibition initiate excessive stress-induced alcohol consumption in mice. PLoS One, 2012, 7:e50187.
269. Kapfhamer D, Taylor S, Zou ME, Lim JP, Kharazia V, Heberlein U. Taok2 controls behavioral response to ethanol in mice. Genes Brain and Behavior, 2013, 12:87-97.
270. Lee MR, Ruby CL, Hinton DJ, Choi S, Adams CA, Young Kang N, Choi DS. Striatal adenosine signaling regulates EAAT2 and astrocytic AQP4 expressn and alcohol drinking in mice. Neuropsychopharmacology, 2013, 38:437-445.
271. Zhou Z, Karlsson C, Liang T, Xiong W, Kimura M, Tapocik JD, Yuan Q, Barbier E, Feng A, Flanigan M, Augier E, Enoch MA, Hodgkinson CA, Shen PH, Lovinger DM, Edenberg HJ, Heilig M, Goldman D. Loss of metabotropic glutamate receptor 2 escalates alcohol consumption. Proceedings of the National Academy of Sciences USA, 2013, 110:16963-16968.
272. Bahi A, Tolle V, Fehrentz JA, Brunel L, Martinez J, Tomasetto CL, Karam SM. Ghrelin knockout mice show decreased voluntary alcohol consumption and reduced ethanol-induced conditioned place preference. Peptides, 2013, 43:48-55.
273. López-Jiménez A, Walter NA, Giné E, Santos Á, Echeverry-Alzate V, Bühler KM, Olmos P, Giezendanner S, Moratalla R, Montoliu L, Buck KJ, López-Moreno JA. A spontaneous deletion of α -synuclein is associated with an increase in CB1 mRNA transcript and receptor expression in the hippocampus and amygdala: effects on alcohol consumption. Synapse, 2013, 67:280-289.
274. Rácz I, Markert A, Mauer D, Stoffel-Wagner B, Zimmer A. Long-term ethanol effects on acute stress responses: modulation by dynorphin. Addiction Biology, 2013, 18:678-688.
275. Conti AC, Lowing JL, Susick LL, Bowen SE. Investigation of calcium-stimulated adenylyl cyclases 1 and 8 on toluene and ethanol neurobehavioral actions. Neurotoxicology and Teratology, 2012, 34:481-488.
276. Dawson A, Miles MF, Damaj MI. The β 2 nicotinic acetylcholine receptor subunit differentially influences ethanol behavioral effects in the mouse. Alcohol, 2013, 47:85-94.
277. Houchi H, Persyn W, Legastelois R, Naassila M. The adenosine A2A receptor agonist CGS 21680 decreases ethanol self-administration in both non-dependent and dependent animals. Addiction Biology, 2013, 18:812-825.

278. Imai S, Kano M, Nonoyama K, Ebihara S. Behavioral characteristics of ubiquitin-specific peptidase 46-deficient mice. PLoS One, 2013, 8:e58566.
279. Nam HW, Hinton DJ, Kang NY, Kim T, Lee MR, Oliveros A, Adams C, Ruby CL, Choi DS. Adenosine transporter ENT1 regulates the acquisition of goal-directed behavior and ethanol drinking through A2A receptor in the dorsomedial striatum. Journal of Neuroscience, 2013, 33:4329-4338.
280. Ruby CL, Walker DL, An J, Kim J, Choi DS. Sex-Specific Regulation of Depression, Anxiety-Like Behaviors and Alcohol Drinking in Mice Lacking ENT1. Journal of Addiction Research and Therapy, 2011, Suppl. 4. doi:pii: 004.
281. Delis F, Thanos PK, Rombola C, Rosko L, Grandy D, Wang GJ, Volkow ND. Chronic mild stress increases alcohol intake in mice with low dopamine D2 receptor levels. Behavioral Neuroscience, 2013, 127:95-105.
282. Lynch LJ, Sullivan KA, Vallender EJ, Rowlett JK, Platt DM, Miller GM. Trace amine associated receptor 1 modulates behavioral effects of ethanol. Substance Abuse, 2013, 7:117-126.
283. Li H, Tao Y, Ma L, Liu X, Ma L. β -Arrestin-2 inhibits preference for alcohol in mice and suppresses Akt signaling in the dorsal striatum. Neuroscience Bulletin, 2013, 29:531-540.
284. Subbanna S, Shivakumar M, Psychoyos D, Xie S, Basavarajappa BS. Anandamide-CB1 receptor signaling contributes to postnatal ethanol-induced neonatal neurodegeneration, adult synaptic, and memory deficits. Journal of Neuroscience, 2013, 33:6350-6366.
285. Stacey D, Bilbao A, Maroteaux M, Jia T, Easton AC, Longueville S, Nymberg C, Banaschewski T, Barker GJ, Büchel C, Carvalho F, Conrod PJ, Desrivières S, Fauth-Bühler M, Fernandez-Medarde A, Flor H, Gallinat J, Garavan H, Bokde AL, Heinz A, Ittermann B, Lathrop M, Lawrence C, Loth E, Lourdasamy A, Mann KF, Martinot JL, Nees F, Palkovits M, Paus T, Pausova Z, Rietschel M, Ruggeri B, Santos E, Smolka MN, Staehlin O, Jarvelin MR, Elliott P, Sommer WH, Mamedi M, Müller CP, Spanagel R, Girault JA, Schumann G; IMAGEN Consortium. RASGRF2 regulates alcohol-induced reinforcement by influencing mesolimbic dopamine neuron activity and dopamine release. Proceedings of the National Academy of Sciences USA, 2012, 109:21128-21133.
286. Santos N, Chatterjee S, Henry A, Holgate J, Bartlett SE. The $\alpha 5$ neuronal nicotinic acetylcholine receptor subunit plays an important role in the sedative effects of ethanol but does not modulate consumption in mice. Alcoholism: Clinical and Experimental Research, 2013, 37:655-662.
287. Giardino WJ, Ryabinin AE. CRF1 receptor signaling regulates food and fluid intake in the drinking-in-the-dark model of binge alcohol consumption. Alcoholism: Clinical and Experimental Research, 2013, 37:1161-1170.

288. Blednov YA, Benavidez JM, Black M, Leiter CR, Osterndorff-Kahanek E, Johnson D, Borghese CM, Hanrahan JR, Johnston GA, Chebib M, Harris RA. GABA_A receptors containing $\rho 1$ subunits contribute to in vivo effects of ethanol in mice. PLoS One, 2014, 9:e85525.
289. Ou XM, Udemgba C, Wang N, Dai X, Lomber G, Seo S, Urrutia R, Wang J, Duncan J, Harris S, Fairbanks CA, Zhang X. Diabetes-causing gene, kruppel-like factor 11, modulates the antinociceptive response of chronic ethanol intake. Alcoholism: Clinical and Experimental Research, 2014, 38:401-408.
290. Kreifeldt M, Le D, Treisman SN, Koob GF, Contet C. BK channel $\beta 1$ and $\beta 4$ auxiliary subunits exert opposite influences on escalated ethanol drinking in dependent mice. Frontiers in Integrative Neuroscience, 2013, 7:105.
291. McCall NM, Sprow GM, Delpire E, Thiele TE, Kash TL, Pleil KE. Effects of sex and deletion of neuropeptide Y2 receptors from GABAergic neurons on affective and alcohol drinking behaviors in mice. Frontiers in Integrative Neuroscience, 2013, 7:100.
292. Wyatt LR, Finn DA, Khoja S, Yardley MM, Asatryan L, Alkana RL, Davies DL. Contribution of P2X4 receptors to ethanol intake in male C57BL/6 mice. Neurochemical Research, 2014, 39:1127-1139.
293. Tseng A, Nguyen K, Hamid A, Garg M, Marquez P, Lutfy K. The role of endogenous beta-endorphin and enkephalins in ethanol reward. Neuropharmacology, 2013, 73:290-300.
294. Susick LL, Lowing JL, Bosse KE, Hildebrandt CC, Chrumka AC, Conti AC. Adenylyl cyclases 1 and 8 mediate select striatal-dependent behaviors and sensitivity to ethanol stimulation in the adolescent period following acute neonatal ethanol exposure. Behavioural Brain Research, 2014, 269:66-74.
295. Salinas AG, Nguyen CTQ, Ahmadi-Tehrani D, Morrisett RA. Reduced ethanol consumption and preference in cocaine- and amphetamine-regulated transcript (CART) knockout mice. Addiction Biology, 2012, 19:175-184.
296. Rial D, Pandolfo P, Bitencourt RM, Pamplona FA, Moreira KM, Hipolide D, Dombrowski PA, Da Cunha C, Walz R, Cunha RA, Takahashi RN, Prediger RD. Cellular prion protein (PrP^c) modulates ethanol-induced behavioral adaptive changes in mice. Behavioural Brain Research, 2014, 271:325-332.
297. Olney JJ, Sprow GM, Navarro M, Thiele TE. The protective effects of the melanocortin receptor (MCR) agonist, melanotan-II (MTII), against binge-like ethanol drinking are facilitated by deletion of the MC3 receptor in mice. Neuropeptides, 2014, 48:47-51.

298. Leggio GM, Camilleri G, Platania CBM, Castorina A, MARRAZZO G, Torrisi SA, Nona CN, D'Agata Y, Nobrega J, Stark H, Bucolo C, Le Foll B, Drago F, Salomone S. Dopamine D3 receptor is necessary for ethanol consumption: an approach with bupirone. Neuropsychopharmacology, 2014, 39:2017-2028.
299. Lee AM, Zou ME, Lim JP, Stecher J, McMahon T, Messing RO. Deletion of Prkcz increases intermittent ethanol consumption in mice. Alcoholism: Clinical and Experimental Research, 2014, 38:170-178.
300. Lamb RJ, Daws LC. Ethanol self-administration in serotonin transporter knockout mice: unconstrained demand and elasticity. Genes, Brain and Behavior, 2013, 12:741-747.
301. Ishiguro H, Hall FS, Horiuchi Y, Sakurai T, Hishimoto A, Grumet M, Uhl GR, Onaivi ES, Arinami T. NrCAM-regulating neural systems and addiction-related behaviors. Addiction Biology, 2012, 19:343-353.
302. Carlson SL, O'Buckley TK, Thomas R, Thiele TE, Morrow AL. Altered GABAA receptor expression and seizure threshold following acute ethanol challenge in mice lacking the RII β subunit of PKA. Neurochemical Research, 2014, 39:1079-1087.
303. Bjork K, Tronci V, Thorsell A, Tanda G, Hirth N, Heilig M, Hansson AC, Sommer WH. β -arrestin 2 knockout mice exhibit sensitized dopamine release and increased reward in response to a low dose of alcohol. Psychopharmacology, 2013, 230:439-449.
304. Liu L, Hendrickson LM, Guildford MJ, Zhao-Shea R, Gardner PD, Tapper AR. Nicotinic acetylcholine receptors containing the α 4 subunit modulate alcohol reward. Biological Psychiatry, 2013, 73:738-746.
305. Ruby CL, Vadnie CA, Hinton DJ, Abulseoud OA, Walker DL, O'Connor KM, Noterman MF, Choi DS. Adenosinergic regulation of striatal clock gene expression and ethanol intake during constant light. Neuropsychopharmacology, 2014, 39:2432-2440.
306. Sachs BD, Salah AA, Caron MG. Congenital brain serotonin deficiency leads to reduced ethanol sensitivity and increased ethanol consumption in mice. Neuropharmacology, 2014, 77:177-184.
307. Vanhanen J, Nuutinen S, Lintunen M, Maki T, Ramo J, Karlstedt K, Panula P. Histamine is required for H3 receptor-mediated alcohol reward inhibition, but not for alcohol consumption or stimulation. British Journal of Pharmacology, 2013, 170:177-187.
308. Cozzoli DK, Courson J, Wroten MG, Greentree DI, Lum EN, Campbell RR, Thompson AB, Maliniak D, Worley PF, Jonquieres G, Klugmann M, Finn DA, Szumlinski KK. Binge alcohol drinking by mice requires intact group I metabotropic glutamate receptor signaling within the central nucleus of the amygdala. Neuropsychopharmacology, 2014, 39:435-444.

309. Herman MA, Sidhu H, Stouffer DG, Kreifeldt M, Le D, Cates-Gatto C, Munoz MB, Roberts AJ, Parsons LH, Roberto M, Wickman K, Slesinger PA, Contet C. GIRK3 gates activation of the mesolimbic dopaminergic pathway by ethanol. Proceedings of the National Academy of Sciences USA, 2015, 112:7091-7096.
310. de la Cour C, Sørensen G, Wortwein G, Weikop P, Dencker D, Fink-Jensen A, Molander A. Enhanced self-administration of alcohol in muscarinic acetylcholine M4 receptor knockout mice. European Journal of Pharmacology, 2015, 746:1-5.
311. Lum EN, Campbell RR, Rostock C, Szumlinski KK. mGluR1 within the nucleus accumbens regulates alcohol intake in mice under limited-access conditions. Neuropharmacology, 2014, 79:679-687.
312. Milner LC, Shirley RL, Kozell LB, Walter NA, Kruse LC, Komiyama NH, Grant SG, Buck KJ. Novel MPDZ/MUPP1 transgenic and knockdown models confirm Mpdz's role in ethanol withdrawal and support its role in voluntary ethanol consumption. Addiction Biology, 2015, 20:143-147.
313. Ortega-Álvaro A, Ternianov A, Aracil-Fernández A, Navarrete F, García-Gutiérrez MS, Manzanares J. Role of cannabinoid CB2 receptor in the reinforcing actions of ethanol. Addiction Biology, 2015, 20:43-55.
314. Savarese A, Zou ME, Kharazia V, Maiya R, Lasek AW. Increased behavioral responses to ethanol in Lmo3 knockout mice. Genes Brain and Behavior, 2014, 13:777-783.
315. Pena-Oliver Y, Sanchez-Roige S, Stephens DN, Ripley TL. Alpha-synuclein deletion decreases motor impulsivity but does not affect risky decision making in a mouse gambling task. Psychopharmacology, 2014, 231:2493-2506.
316. Vicente-Rodriguez M, Perez-Garcia C, Ferrer-Alcon M, Uribarri M, Sanchez-Alonso MG, Ramos MP, Herradon G. Pleiotrophin differentially regulates the rewarding and sedative effects of ethanol. Journal of Neurochemistry, 2014, 131:688-695.
317. Stewart A, Maity B, Andereg SP, Allamargot C, Yang J, Fisher RA. Regulator of G protein signaling 6 is a critical mediator of both reward-related behavioral and pathological responses to alcohol. Proceedings of the National Academy of Sciences USA, 2015, 112: E786-E795.
318. Huang MC, Schwandt ML, Chester JA, Kirchhoff AM, Kao CF, Liang T, Tapocik JD, Ramchandani VA, George DT, Hodgkinson CA, Goldman D, Heilig M. FKBP5 moderates alcohol withdrawal severity: human genetic association and functional validation in knockout mice. Neuropsychopharmacology, 2014, 39:2029-2038.

319. Walker AW, Smith CM, Chua BE, Krstew EV, Zhang C, Gundlach AL, Lawrence AJ. Relaxin-3 receptor (RXFP3) signalling mediates stress-related alcohol preference in mice. PLoS One, 2015, 10:e0122504.
320. Raud S, Reimets R, Loomets M, Sutt S, Altpere A, Visnapuu T, Innos J, Luuk H, Plaas M, Volke V, Vasar E. Deletion of the Wolfram syndrome-related gene *Wfs1* results in increased sensitivity to ethanol in female mice. Neuropharmacology, 2015, 95:59-67.
321. Legastelois R, Darcq E, Wegner SA, Lombroso PJ, Ron D. Striatal-enriched protein tyrosine phosphatase controls responses to aversive stimuli: implication for ethanol drinking. PLoS One, 2015, 10:e0127408.
322. Vicente-Rodriguez M, Perez-Garcia C, Haro M, Ramos MP, Herradon G. Genetic inactivation of midkine modulates behavioural responses to ethanol possibly by enhancing GABA(A) receptor sensitivity to GABA(A) acting drugs. Behavioural Brain Research, 2014, 274:258-263.
323. Contet C, Kim A, Le D, Iyengar SK, Kotzebue RW, Yuan CJ, Kieffer BL, Mandyam CD. μ -Opioid receptors mediate the effects of chronic ethanol binge drinking on the hippocampal neurogenic niche. Addiction Biology, 2014, 19:770-780.
324. Blednov YA, Benavidez JM, Black M, Leiter CR, Osterndorff-Kahanek E, Harris RA. Glycine receptors containing $\alpha 2$ or $\alpha 3$ subunits regulate specific ethanol-mediated behaviors. Journal of Pharmacology and Experimental Therapeutics, 2015, 353:181-191.
325. Guildford MJ, Sacino AV, Tapper AR. Modulation of ethanol reward sensitivity by nicotinic acetylcholine receptors containing the $\alpha 6$ subunit. Alcohol, 2016, 57:65-70.
326. Ford MM, Nickel JD, Kaufman MN, Finn DA. Null mutation of 5 α -reductase type I gene alters ethanol consumption patterns in a sex-dependent manner. Behavior Genetics, 2015, 45:341-353.
327. Gavin DP, Kusumo H, Zhang H, Guidotti A, Pandey SC. Role of growth arrest and DNA damage-inducible, beta in alcohol-drinking behaviors. Alcoholism: Clinical and Experimental Research, 2016, 40:263-272.
328. Reker AN, Oliveros A, Sullivan JM 3rd, Nahar L, Hinton DJ, Kim T, Bruner RC, Choi DS, Goeders NE, Nam HW. Neurogranin in the nucleus accumbens regulates NMDA receptor tolerance and motivation for ethanol seeking. Neuropharmacology, 2018, 131:58-67.
329. Blasio A, Wang J, Wang D, Varodayan FP, Pomrenze MB, Miller J, Lee AM, McMahon T, Gyawali S, Wang HY, Roberto M, McHardy S, Pleiss MA, Messing RO. Novel small-molecule inhibitors of protein kinase C epsilon reduce ethanol consumption in mice. Biological Psychiatry, 2018, 84:193-201.

330. Haass-Koffler CL, Henry AT, Melkus G, Simms JA, Naemmuddin M, Nielsen CK, Lasek AW, Magill M, Schwandt ML, Momenan R, HODgkinson CA, Bartlett SE, Swift RM, Bonci A, Leggio L. Defining the role of corticotropin releasing factor binding protein in alcohol consumption. Translational Psychiatry, 2016, 6:e953.
331. Powers MS, Breit KR, Chester JA. Genetic versus pharmacological assessment of the role of cannabinoid type 2 receptors in alcohol reward-related behaviors. Alcoholism: Clinical and Experimental Research, 2015, 39:2438-2446.
332. Patkar OL, Belmer A, Tarren JR, Holgate JY, Bartlett SE. The effect of varenicline on binge-like ethanol consumption in mice is $\beta 4$ nicotinic acetylcholine receptor-independent. Neuroscience Letters, 2016, 633:235-239.
333. Van't Veer A, Smnith KL, Cohen BM, Carlezon WA Jr, Bechtholt AJ. Kappa-opioid receptors differentially regulate low and high levels of ethanol intake in female mice. Brain and Behavior, 2016, 6:e00523.
334. Qiu B, Luczak SE, Wall TL, Kirchoff AM, Xu Y, Eng MY, Stewart RB, Shou W, Boehm SL 2nd, Chester JA, Yong W, Liang T. The *FKBP5* gene affects alcohol drinking in knockout mice and is implicated in alcohol drinking in humans. International Journal of Molecular Sciences, 2016, 17:1271.
335. McGonigle CE, Nentwig TB, Wilson DE, Rhinehart EM, Grisel JE. β -endorphin regulates alcohol consumption induced by exercise restriction in female mice. Alcohol, 2016, 53:51-60.
336. Schweitzer P, Cates-Gatto C, Varodayan FP, Nadav T, Roberto M, Lasek AW, Roberts AJ. Dependence-induced ethanol drinking and GABA neurotransmission are altered in *Alk* deficiency mice. Neuropharmacology, 2016, 107:1-8.
337. Jang Y, Lee MH, Park JH, Han SY, Kim DK. TLR3 deficiency increases voluntary alcohol consumption. Neuroreport, 2016, 27:356-360.
338. Blednov YA, Black M, Benavidez JM, Stamatakis EE, Harris RA. PPAR agonists: I. Role of receptor subunits in alcohol consumption in male and female mice. Alcoholism: Clinical and Experimental Research, 2016, 40:553-562.
339. Shirahase T, Aoki M, Watanabe R, Watanabe Y, Tanaka M. Increased alcohol consumption in relaxin-3 deficient male mice. Neuroscience Letters, 2016, 612:155-160.
340. Valenza M, DiLeo A, Steardo L, Cottone P, Sabino V. Ethanol-related behaviors in mice lacking the sigma-1 receptor. Behavioural Brain Research, 2016, 297:196-203.

341. Walker AW, Smith CM, Gundlach AL, Lawrence AJ. Relaxin-3 receptor (Rxfp3) gene deletion reduces operant sucrose- but not alcohol-responding in mice. Genes, Brain and Behavior, 2015, 14:625-634.
342. Blednov YA, Benavidez JM, Black M, Mayfield J, Harris RA. Role of interleukin-1 receptor signaling in the behavioral effects of ethanol and benzodiazepines. Neuropharmacology, 2015, 95:309-320.
343. Pradier B, Erxlebe E, Markert A, Racz I. Interaction of cannabinoid receptor 2 and social environment modulates chronic alcohol consumption. Behavioural Brain Research, 2015, 287:163-171.
344. Moriya Y, Kasahara Y, Hall FS, Sakakibara Y, Uhl GR, Tomita H, Sora I. Sex differences in the effects of adolescent social deprivation on alcohol consumption in μ -opioid receptor knockout mice. Psychopharmacology, 2015, 232:1471-1482.
345. Smith ML, Li J, Ryabinin AE. Increased alcohol consumption in urocortin 3 knockout mice is unaffected by chronic inflammatory pain. Alcohol and Alcoholism, 2015, 50:132-139.
346. Shannon EE, Shelton KL, Vivian JA, Yount I, Morgan AR, Homanics GE, Grant KA. Discriminative stimulus effects of ethanol in mice lacking the γ -aminobutyric acid type A receptor δ subunit. Alcoholism: Clinical and Experimental Research, 2004, 28:906-913.
347. Chen H, He D, Lasek AW. Midkine in the mouse ventral tegmental area limits ethanol intake and Ccl2 gene expression. Genes, Brain and Behavior, 2017, 16:699-708.
348. Schumann G, Liu C, O'Reilly P, Gao H, Song P, Xu B, Ruggeri B, Amin N, Jia T, Preis S, Segura Lepe M, Akira S, Barbieri C, Baumeister S, Cauchi S, Clarke TK, Enroth S, Fischer K, Hällfors J, Harris SE, Hieber S, Hofer E, Hottenga JJ, Johansson Å, Joshi PK, Kaartinen N, Laitinen J, Lemaitre R, Loukola A, Luan J, Lyytikäinen LP, Mangino M, Manichaikul A, Mbarek H, Milanesechi Y, Moayyeri A, Mukamal K, Nelson C, Nettleton J, Partinen E, Rawal R, Robino A, Rose L, Sala C, Satoh T, Schmidt R, Schraut K, Scott R, Smith AV, Starr JM, Teumer A, Trompet S, Uitterlinden AG, Venturini C, Vergnaud AC, Verweij N, Vitart V, Vuckovic D, Wedenoja J, Yengo L, Yu B, Zhang W, Zhao JH, Boomsma DI, Chambers J, Chasman DI, Daniela T, de Geus E, Deary I, Eriksson JG, Esko T, Eulenburger V, Franco OH, Froguel P, Gieger C, Grabe HJ, Gudnason V, Gyllenstein U, Harris TB, Hartikainen AL, Heath AC, Hocking L, Hofman A, Huth C, Jarvelin MR, Jukema JW, Kaprio J, Kooner JS, Kutalik Z, Lahti J, Langenberg C, Lehtimäki T, Liu Y, Madden PA, Martin N, Morrison A, Penninx B, Pirastu N, Psaty B, Raitakari O, Ridker P, Rose R, Rotter JI, Samani NJ, Schmidt H, Spector TD, Stott D, Strachan D, Tzoulaki I, van der Harst P, van Duijn CM, Marques-Vidal P, Vollenweider P, Wareham NJ, Whitfield JB, Wilson J, Wolffenbuttel B, Bakalkin G, Evangelou E, Liu Y, Rice KM, Desrivieres S, Kliewer SA, Mangelsdorf DJ, Müller CP, Levy D, Elliott P. KLB is associated with alcohol drinking, and its gene product β -Klotho is necessary for

- FGF21 regulation of alcohol preference. Proceedings of the National Academy of Sciences USA, 2016, 113:14372-14377.
349. Robins MT, Chiang T, Berry JN, Ko MJ, Ha JE, van Rijn RM. Behavioral characterization of β -arrestin 1 knockout mice in anxiety-like and alcohol behaviors. Frontiers in Behavioral Neuroscience, 2018, 12:54.
 350. Germany CE, Reker AN, Hinton DJ, Oliveros A, Shen X, Andres-Beck LG, Winger KM, Trutschl M, Cvek U, Choi DS, Nam HW. Pharmacoproteomics profile in response to acamprosate treatment of an alcoholism animal model. Proteomics, 2018, 18:1700417.
 351. Chiang T, Sansuk K, van Rijn RM. β -arrestin 2 dependence of δ opioid receptor agonists is correlated with alcohol intake. British Journal of Pharmacology, 2016, 173:332-343.
 352. Blednov YA, Black M, Chernis J, Da Costa A, Mayfield J, Harris RA. Ethanol consumption in mice lacking CD14, TLR2, TLR4, or MyD88. Alcoholism: Clinical and Experimental Research, 2017, 41:516-530.
 353. Castilla-Ortega E, Pavon FJ, Sanchez-Marin L, Estivill-Torres G, Pedraza C, Blanco E, Suarez J, Santin L, Rodriguez de Fonseca F, Serrano A. Both genetic deletion and pharmacological blockade of lysophosphatidic acid LPA1 receptor results in increased alcohol consumption. Neuropharmacology, 2016, 103:92-103.
 354. Stefaniuk M, Beroun A, Lebitko T, Markina O, Leski S, Meyza K, Grzywacz A, Samochowiec J, Samochowiec A, Radwanska K, Kaczmarek L. Matrix metalloproteinase-9 and synaptic plasticity in the central amygdala in control of alcohol-seeking behavior. Biological Psychiatry, 2017, 81:907-917.
 355. Kamens HM, Silva C, McCarthy R, Cox RJ, Ehringer MA. No evidence of a role of the β 4 subunit of the nicotinic acetylcholine receptor in alcohol-related behaviors. BMC Research Notes, 2017, 10:151.
 356. Sumitomo A, Ueta K, Mauchi S, Hirai K, Horike K, Hikida T, Sakurai T, Sawa A, Tomoda T. Ulk1 protects against ethanol-induced neuronal stress and cognition-related behavioral deficits. Neuroscience Research, 2017, 117:54-61.
 357. Repunte-Canonigo V, Herman MA, Kawamura T, Kranzler HR, Sherva R, Gelernter J, Farrer LA, Roberto M, Sanna PP. Nf1 regulates alcohol dependence-associated excessive drinking and gamma-aminobutyric acid release in the central amygdala in mice and is associated with alcohol dependence in humans. Biological Psychiatry, 2015, 77:870-879.
 358. Karlsson C, Schank JR, Rehman F, Stojakovic A, Bjork K, Barbier E, Solomon M, Tapocik J, Engblom D, Thorsell A, Heilig M. Proinflammatory signaling regulates voluntary alcohol intake and stress-induced consumption after exposure to social defeat stress in mice. Addiction Biology, 2016, 22:1279-1288.

359. Montesinos J, Gil A, Guerri C. Nalmefene prevents alcohol-induced neuroinflammation and alcohol drinking preference in adolescent female mice: role of TLR4. Alcoholism: Clinical and Experimental Research, 2017, 41, 1257-1270.
360. Garcia-Gutierrez MS, Navarrete F, Laborda J, Manzanares J. Deletion of Dlk1 increases the vulnerability to developing anxiety-like behaviors and ethanol consumption in mice. Biochemical Pharmacology, 2018, 158:37-44.
361. McMurray KMJ, Sidhu PS, Cook JM, Arnold LA, Palmer AA. Genetic and pharmacological manipulation of glyoxalase 1 regulates voluntary ethanol consumption in mice. Addiction Biology, 2015, 22:381-389.
362. Montesinos J, Pascual M, Rodriguez-Arias M, Minarro J, Guerri C. Involvement of TLR4 in the long-term epigenetic changes, rewarding and anxiety effects induced by intermittent ethanol treatment in adolescence. Brain, Behavior, and Immunity, 2016, 53:159-171.
363. Kozell LB, Denmark DL, Walter NAR, Buck KJ. Distinct roles for two chromosome 1 loci in ethanol withdrawal, consumption, and conditioned place preference. Frontiers in Genetics, 2018, 9:323.
364. Khoja S, Kuynh N, Asatryan L, Jakowec MW, Davies DL. Reduced expression of purinergic P2X4 receptors increases voluntary ethanol intake in C57BL/6J mice. Alcohol, 2018, 68:63-70.
365. Jury NJ, Radke AK, Pati D, Kocharian A, Mishina M, Kash TL, Holmes A. NMDA receptor GluN2A subunit deletion protects against dependence-like ethanol drinking. Behavioural Brain Research, 2018, 353:124-128.
366. Wooden JI, Schuller K, Roman G, Das J, Leasure JL. Munc 13-1 heterozygosity does not alter voluntary ethanol consumption or sensitivity in mice. Alcohol, 2019, in press.
367. Bosse KE, Ghoddoussi F, Eapen AT, Charlton JL, Susick LL, Desai K, Berkowitz BA, Perrine SA, Conti AC. Calcium/calmodulin-stimulated adenylyl cyclases 1 and 8 regulate reward-related brain activity and ethanol consumption. Brain Imaging and Behavior, 2019, 13:396-407.
368. Jia YF, Vadnie CA, Ho AMC, Peyton L, Veldic M, Wininger K, Matveyenko A, Choi DS. Type 1 equilibrative nucleoside transporter (ENT1) regulates sex-specific ethanol drinking during disruption of circadian rhythms. Addiction Biology, 2019, e12801.
369. Olsen D, Kaas M, Lundhede J, Molgaard S, Sykjaer A, Kjolby M, Ostergaard SD, Glerup S. Reduced alcohol seeking and withdrawal symptoms in mice lacking the BDNF receptor SorCS2. Frontiers in Pharmacology, 10:499.

370. Kalinichenko LS, Muhle C, Eulenburg V, Praetner M, Reichel M, Gulbins E, Kornhuber J, Muller CP. Enhanced alcohol preference and anxiolytic alcohol effects in Niemann-Pick disease model in mice. Frontiers in Neurology, 2019, 10:731.
371. Ben Hamida S, Mendonça-Netto S, Arefin TM, Nasseef MT, Boulos LJ, McNicholas M, Ehrlich AT, Clarke E, Moquin L, Gratton A, Darcq E, Harsan LA, Maldonado R, Kieffer BL. Increased alcohol seeking in mice lacking Gpr88 involves dysfunctional mesocorticolimbic networks. Biological Psychiatry, 2018, 84:202-212.
372. Brunner SM, Farzi A, Locker F, Holub BS, Drexel M, Reichmann F, Lang AA, Mayr JA, Vilches JJ, Navarro X, Lang R, Sperk G, Holzer P, Kofler B. GAL3 receptor KO mice exhibit an anxiety-like phenotype. Proceedings of the National Academy of Sciences USA, 2018, 24:886-897.
373. Robins MT, Chiang T, Mores KL, Alongkronrusmee D, van Rijn RM. Critical role for G(i/o)-protein activity in the dorsal striatum in the reduction of voluntary alcohol intake in C57Bl/6 mice. Frontiers in Psychiatry, 2018, 9:112.
374. Mielenz D, Reichel M, Jia T, Quinlan EB, Stöckl T, Mettang M, Zilske D, Kirmizi-Alsan E, Schönberger P, Praetner M, Huber SE, Amato D, Schwarz M, Purohit P, Brachs S, Spranger J, Hess A, Büttner C, Ekici AB, Perez-Branguli F, Winner B, Rauschenberger V, Banaschewski T, Bokde ALW, Büchel C, Conrod PJ, Desrivières S, Flor H, Frouin V, Gallinat J, Garavan H, Gowland P, Heinz A, Martinot JL, Lemaitre H, Nees F, Paus T, Smolka MN; IMAGEN Consortium, Schambony A, Bäumler T, Eulenburg V, Alzheimer C, Lourdasamy A, Schumann G, Müller CP. EFhd2/Swiprosin-1 is a common genetic determinant for sensation-seeking/low anxiety and alcohol addiction. Molecular Psychiatry, 2018, 23:1303-1319.
375. Leggio GM, Di Marco R, Gulisano W, D'Ascenzo M, Torrisi SA, Geraci F, Lavanco G, Dahl K, Giurdanella G, Castorina A, Aitta-Aho T, Aceto G, Bucolo C, Puzzo D, Grassi C, Korpi ER, Drago F, Salomone S. Dopaminergic-GABAergic interplay and alcohol binge drinking. Pharmacological Research, 2019, 141:384-391.
376. Schank JR, Lee S, Gonzalez-Islas CE, Nennig SE, Fulenwider HD, Chang J, Li JM, Kim Y, Jeffers LA, Chung J, Lee JK, Jin Z, Aalkjaer C, Boedtkjer E, Choi I. Increased alcohol consumption in mice lacking sodium bicarbonate transporter NBCn1. Sci Rep 2020;10(1):11017.
377. Blednov YA, Da Costa A, Mayfield J, Harris RA, Messing RO. Deletion of Tlr3 reduces acute tolerance to alcohol and alcohol consumption in the intermittent access procedure in male mice. Addict Biol 2021;26(2):e12932.
378. Rodriguez KM, Smith BL, Caldwell HK. Voluntary alcohol consumption is increased in female, but not male, oxytocin receptor knockout mice. Brain Behav 2020;10(9):e01749.

379. Müller CP, Chu C, Qin L, Liu C, Xu B, Gao H, Ruggeri B, Hieber S, Schneider J, Jia T, Tay N, Akira S, Satoh T, Banaschewski T, Bokde ALW, Bromberg U, Büchel C, Quinlan EB, Flor H, Frouin V, Garavan H, Gowland P, Heinz A, Ittermann B, Martinot JL, Martinot MP, Artiges E, Lemaitre H, Nees F, Papadopoulos Orfanos D, Paus T, Poustka L, Millenet S, Fröhner JH, Smolka MN, Walter H, Whelan R, Bakalkin G, Liu Y, Desrivières S, Elliott P, Eulenburg V, Levy D, Crews F, Schumann G. The cortical neuroimmune regulator TANK affects emotional processing and enhances alcohol drinking: a translational study. *Cereb Cortex* 2019;29(4):1736-1751.
380. Nentwig TB, Wilson DE, Rhinehart EM, Grisel JE. Sex differences in binge-like EtOH drinking, corticotropin-releasing hormone and corticosterone: effects of β -endorphin. *Addict Biol* 2019;24(3):447-457.
381. Ben Hamida S, Boulos LJ, McNicholas M, Charbogne P, Kieffer BL. Mu opioid receptors in GABAergic neurons of the forebrain promote alcohol reward and drinking. *Addict Biol* 2019;24(1):28-39.
382. Guillot A, Ren T, Jourdan T, Pawlosky RJ, Han E, Kim SJ, Zhang L, Koob GF, Gao B. Targeting liver aldehyde dehydrogenase-2 prevents heavy but not moderate alcohol drinking. *Proc Natl Acad Sci U S A* 2019;116(51):25974-25981.
383. Salling MC, Harrison NL. Constitutive genetic deletion of *Hcn1* increases alcohol preference during adolescence. *Brain Sci* 2020;10(11):763.
384. San Martin L, Gallegos S, Araya A, Romero N, Morelli G, Comhair J, Harvey RJ, Rigo JM, Brone B, Aguayo LG. Ethanol consumption and sedation are altered in mice lacking the glycine receptor $\alpha 2$ subunit. *Br J Pharmacol* 2020;177(17):3941-3956.